

# Isolated Horizons and Black Hole Entropy In Loop Quantum Gravity

Jacobo Diaz-Polo<sup>\*1,2</sup> and Daniele Pranzetti<sup>†3</sup>

<sup>1</sup>*Department of Physics and Astronomy, Louisiana State University,  
Baton Rouge, LA 70803-4001, USA*

<sup>2</sup>*Institute for Gravitation and the Cosmos & Physics Department,  
Penn State, University Park, PA 16802-6300, USA*

<sup>3</sup>*Max Planck Institute for Gravitational Physics (AEI), Am  
Mühlenberg 1, D-14476 Golm, Germany*

December 2, 2011

## Abstract

We review the black hole entropy calculation in the framework of Loop Quantum Gravity based on the quasi-local definition of a black hole encoded in the isolated horizon formalism. We show, by means of the covariant phase space formalism, the appearance in the conserved symplectic structure of a boundary term corresponding to a Chern-Simons theory on the horizon and present its quantization both in the  $U(1)$  gauge fixed version and in the fully  $SU(2)$  invariant one. We then describe the boundary degrees of freedom counting techniques developed for an infinite value of the Chern-Simons level case and, less rigorously, for the case of a finite value. This allows us to perform a comparison between the  $U(1)$  and  $SU(2)$  approaches and provide a state of the art analysis of their common features and different implications for the entropy calculations. In particular, we comment on different points of view regarding the nature of the horizon degrees of freedom and the role played by the Barbero-Immirzi parameter. We conclude by presenting some of the most recent results concerning possible observational tests for theory.

---

\*jacobop@phys.lsu.edu

†pranzetti@aei.mpg.de

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
<b>2</b>	<b>Definition of Isolated Horizons</b>	<b>8</b>
2.1	IH Classification According to Their Symmetry Groups . . . . .	16
2.2	IH Classification According to the Reality of $\Psi_2$ . . . . .	17
2.3	The Horizon Constraints . . . . .	17
<b>3</b>	<b>The Conserved Symplectic Structure</b>	<b>20</b>
3.1	Action Principle and Phase-Space . . . . .	20
3.2	The Conserved Symplectic Structure in Terms of Vector Variables . . . . .	22
3.3	The Conserved Symplectic Structure in Terms of Real Connection Variables . . . . .	23
3.3.1	The Spherically Symmetric Case . . . . .	25
<b>4</b>	<b>The Zeroth and First Laws of BH Mechanics for Isolated Horizons</b>	<b>27</b>
<b>5</b>	<b>Quantization</b>	<b>28</b>
5.1	U(1) Quantization . . . . .	29
5.1.1	Bulk Hilbert Space . . . . .	29
5.1.2	Surface Hilbert Space . . . . .	31
5.1.3	Quantum Boundary Conditions . . . . .	33
5.2	SU(2) Quantization . . . . .	35
<b>6</b>	<b>Entropy Computation</b>	<b>40</b>
6.1	The Infinite $k$ Counting . . . . .	43
6.1.1	Area Spectrum Characterization . . . . .	45
6.1.2	Generating Functions . . . . .	48
6.1.3	Computational Implementation and Analysis of the Results . . . . .	48
6.1.4	Large Area Asymptotics . . . . .	52
6.2	The Finite $k$ Counting . . . . .	53
<b>7</b>	<b>On the Nature of the Entropy Degrees of Freedom</b>	<b>56</b>
<b>8</b>	<b>On the Role of the Barbero-Immirzi Parameter</b>	<b>60</b>
<b>9</b>	<b>Derivation from Conformal Field Theories</b>	<b>63</b>
<b>10</b>	<b>Observational Tests</b>	<b>66</b>
<b>11</b>	<b>Conclusions</b>	<b>69</b>

# 1 Introduction

Black holes are intriguing solutions of classical general relativity describing important aspects of the physics of gravitational collapse. Their existence in our nearby universe is by now supported by a great amount of observational evidence [1]. When isolated, these systems are remarkably simple for late and distant observers: once the initial very dynamical phase of collapse is passed the system is expected to settle down to a stationary situation completely described (as implied by the famous results by Carter, Israël, and Hawking [2]) by the three extensive parameters (mass  $M$ , angular momentum  $J$ , electric charge  $Q$ ) of the Kerr-Newman family [3].

However, the great simplicity of the final stage of an isolated gravitational collapse for late and distant observers is in sharp contrast with the very dynamical nature of the physics seen by in-falling observers which depends on all the details of the collapsing matter. Moreover, this dynamics cannot be consistently described for late times (as measured by the in-falling observers) using General Relativity due to the unavoidable development, within the classical framework, of unphysical pathologies of the gravitational field. Concretely, the celebrated singularity theorems of Hawking and Penrose [4] imply the breakdown of predictability of General Relativity in the black hole interior. Dimensional arguments imply that quantum effects cannot be neglected near the classical singularities. Understanding of physics in this extreme regime requires a quantum theory of gravity (see, e.g., [5]-[7]). Black holes (BH) provide, in this precise sense, the most tantalizing theoretical evidence for the need of a more fundamental (quantum) description of the gravitational field.

Extra motivation for the quantum description of gravitational collapse comes from the physics of black holes available to observers outside the horizon. As for the interior physics, the main piece of evidence comes from the classical theory itself which implies an (at first only) apparent relationship between the properties of idealized black hole systems and those of thermodynamical systems. On the one hand, black hole horizons satisfy the very general Hawking area theorem (the so-called *second law*) stating that the black hole horizon area  $a_H$  can only increase, namely

$$\delta a_H \geq 0. \quad (1)$$

On the other hand, the uniqueness of the Kerr-Newman family, as the final (stationary) stage of the gravitational collapse of an isolated gravitational system, can be used to prove the first and zeroth laws: under external perturbation the initially stationary state of a black hole can change but the final stationary state will be described by another Kerr-Newman solution whose parameters readjust according to the *first law*

$$\delta M = \frac{\kappa_H}{8\pi G} \delta a_H + \Phi_H \delta Q + \Omega_H \delta J, \quad (2)$$

where  $\kappa_H$  is the surface gravity,  $\Phi_H$  is the electrostatic potential at the horizon, and  $\Omega_H$  the angular velocity of the horizon. There is also the *zeroth law* stating the uniformity of the surface gravity  $\kappa_H$  on the event horizon of stationary black holes, and finally *the third law* precluding the possibility of reaching an extremal black hole (for which  $\kappa_H = 0$ ) by means of any physical process<sup>1</sup>.

The validity of these classical laws motivated Bekenstein [8] to put forward the idea that black holes may behave as thermodynamical systems with an entropy  $S = \alpha a/\ell_p^2$  and a temperature  $kT = \hbar\kappa_H/(8\pi\alpha)$  where  $\alpha$  is a dimensionless constant and the dimensionality of the quantities involved require the introduction of  $\hbar$  leading in turn to the appearance of the Planck length  $\ell_p$ . The key point is that the need of  $\hbar$  required by the dimensional analysis involved in the argument called for the investigation of black hole systems from a quantum perspective.

In fact, not long after, the semiclassical calculations of Hawking [9] – that studied particle creation in a quantum test field (representing quantum matter and quantum gravitational perturbations) on the space-time background of the gravitational collapse of an isolated system described for late times by a stationary black hole – showed that once black holes have settled to their stationary (classically) final states, they continue to radiate as perfect black bodies at temperature  $kT = \kappa_H\hbar/(2\pi)$ . Thus, on the one hand, this confirmed that black holes are indeed thermal objects that radiate at a given temperature and whose entropy is given by  $S = a/(4\ell_p^2)$ , while, on the other hand, this raised a wide range of new questions whose proper answer requires a quantum treatment of the gravitational degrees of freedom.

Among the simplest questions is the issue of the statistical origin of black hole entropy. In other words, what is the nature of the large amount of micro-states responsible for black hole entropy. This simple question cannot be addressed using semiclassical arguments of the kind leading to Hawking radiation and requires a more fundamental description. In this way, the computation of black hole entropy from basic principles became an important test for any candidate quantum theory of gravity.

In String Theory the entropy has been computed using dualities and no-normalization theorems valid for extremal black holes [10]. There are also calculations based on the effective description of near horizon quantum degrees of freedom in terms of effective 2-dimensional conformal theories [11, 12]. In the rest of this work, we are going to review the quantum description of the microscopic degrees of freedom of a black hole horizon and the derivation of its entropy in the framework of Loop Quantum Gravity (LQG) [13]. In all cases agreement with the Bekenstein-Hawking formula is obtained with logarithmic corrections in  $a/\ell_p^2$ .

---

<sup>1</sup>The third law can only be motivated by a series of examples. Extra motivations come from the validity of the cosmic censorship conjecture.

In LQG, the basic conceptual ideas leading to the black hole entropy calculation date back to the mid nineties and bloomed out of the beautiful interplay between some pioneering works by Smolin, Rovelli and Krasnov.

In [14] Smolin investigated the emergence of the Bekenstein bound and the holographic hypothesis in the context of non-perturbative quantum gravity by studying the quantization of the gravitational field in the case where self-dual boundary conditions are imposed on a boundary with finite spatial area. This was achieved through the construction of an isomorphism between the states and observables of  $SU(2)$  Chern-Simons theory on the boundary and quantum gravity. This correspondence supported the assumption that the space of states of the quantum gravitational field in the bulk region must be spanned by eigenstates of observables that are functions of fields on the boundary and provided the following picture. The metric of a spatial surface turns out to label the different topological quantum field theories that may be defined on it. The physical state space that describes the 4 dimensional quantum gravitational field in a region bounded by that surface will then be constructed from the state spaces of all the topological quantum field theories that live on it.

In [15] Rovelli obtained a black hole entropy proportional to the area by performing computations (valid for physical black holes) based on general considerations and the fact that the area spectrum in the theory is discrete. He suggested that the black hole entropy should be related to the number of quantum microstates of the horizon geometry which correspond to a given macroscopic configuration and are distinguishable from the exterior of the hole.

Combining the main ingredients of these two works then, Krasnov provided [16] a description of the microscopic states of Schwarzschild black hole in terms of states of  $SU(2)$  Chern-Simons theory. Using this description as the basis of a statistical mechanical analysis, he found that the entropy contained within the black hole is proportional to the area of the horizon, with a proportionality coefficient which turns out to be a function of the Barbero-Immirzi parameter.

These fundamental steps provided a solid conceptual (and also technical) basis to the seminal works of [17, 18, 19], which followed right after. Here the authors started from the important observation that the very notion of black hole – as the region causally disconnected from future null infinity – becomes elusive in the context of quantum gravity. This is due to the simple fact that black hole radiation in the semiclassical regime imply that in the full quantum theory the global structure of space-time (expected to make sense away from the strong field region) might completely change – in fact, recent models in two dimensions support the view that this is the case [20] –. For that reason, the problem of black hole entropy in quantum gravity requires the use of a local or quasi-local notion of horizon in equilibrium. Such a local definition of BH has been introduced [21] (see also [22]-[24]) through the

concept of Isolated Horizons (IH). Isolated horizons are regarded as a sector of the phase-space of GR containing a horizon in “equilibrium” with the external matter and gravitational degrees of freedom. This local definition provided a general framework to apply to the black hole entropy calculation in the context of LQG, as first performed (for spherically symmetric IH) in [19]. In this work the authors show, after introduction of a suitable gauge fixing, how the degrees of freedom that are relevant for the entropy calculation can be encoded in a boundary  $U(1)$  Chern-Simons theory.

After separately quantizing the bulk and the boundary theory of the system and imposing the quantum version of the horizon boundary condition, bulk and boundary degrees of freedom are again related to each other and (the ‘gauge fixed’ version of) Smolin’s picture is recovered [19]. By counting the number of states in the boundary Hilbert space, tracing over the bulk degrees, [19] found a leading term for the horizon entropy matching the Bekenstein-Hawking area law, as long as the Barbero-Immirzi parameter  $\beta$  is fixed to a given numerical value. From this point on, the semi-classical result of Bekenstein and Hawking started to be regarded as physical ‘external’ input to fix the ambiguity affecting the non-perturbative quantum theory of geometry.

Soon after this construction of quantum isolated horizons, there has been a blooming of literature devoted to the improvement of the counting problem and which led to the important discoveries of sub-leading logarithmic corrections as well as of a discrete structure of the entropy for small values of the horizon area. First, in [25], a reformulation of the counting problem was performed according to the spirit of the original derivations, and solving certain incompatibilities of the previous computations. An asymptotic computation of entropy, based on this new formulation of the problem, was performed in [26], yielding a first order logarithmic correction to the leading linear behavior. Alternative approaches to the counting and the computation of logarithmic correction were also worked out in [27, 28].

In [29, 30], an exact detailed counting was performed for the first time, showing the discretization of entropy as a function of area for microscopic black holes. Several works followed [31, 32, 33, 34], analyzing these effect from several points of view. A more elegant and technically advanced exact solution, involving analytical methods, number theory, and generating functions was developed in [35, 36, 37], providing the arena for the extension of the exact computation to the large area regime, studied in [38].

However, despite the great enthusiasm fueled by these results, some features of the entropy calculation in LQG were not fully satisfactory. First of all, the need to fix a purely quantum ambiguity represented by the Barbero-Immirzi parameter through the request of agreement with a semi-classical result (coming from a quantum field theory calculation in curved spacetimes with large isolated horizons) didn’t seem a very natural, let alone elegant, passage to many. Moreover, a controversy appeared in the liter-

ature concerning the specific numerical value which  $\beta$  should be tuned to and a discrepancy was found between the constant factor in front of the logarithmic corrections obtained in the  $U(1)$  symmetry reduced model and the seemingly very general treatment (which includes the String Theory calculations [10]) proposed by Carlip [12]. All this stressed the necessity of a more clear-cut relationship between the boundary theory and the LQG quantization of bulk degrees of freedom. Finally, a fully  $SU(2)$  invariant treatment of the horizon degrees appeared more appropriate also from the point of view of the original conceptual considerations of [14, 15, 16, 39].

The criticisms listed above motivated the more recent analysis of [40, 41, 42], which clarified the description of both the classical as well as the quantum theory of black holes in LQG making the full picture more transparent. In fact, these works showed that the gauge symmetry of LQG need not be reduced from  $SU(2)$  to  $U(1)$  at the horizon, leading to a drastic simplification of the quantum theory in which states of a black hole are now in one-to-one correspondence with the fundamental basic volume excitations of LQG given by single intertwiner states. This  $SU(2)$  invariant formulation – equivalent to the  $U(1)$  at the classical level – preserves, in the spherically symmetric case, the Lie algebraic structure of the boundary conditions also at the quantum level, allowing for the proper Dirac imposition of the constraints and the correct restriction of the number of admissible boundary states. In this way, the factor  $-3/2$  in front of the logarithmic corrections is recovered, as shown in [43], eliminating the apparent tension with other approaches to entropy calculation. Moreover, the more generic nature of the  $SU(2)$  treatment allows to loose the numerical restriction on the value of  $\beta$ , by only requiring a given relationship between the parameter in the bulk theory (LQG) and the analog in the boundary theory (the Chern-Simons level), in order to recover the Bekenstein-Hawking entropy [42, 44] – the resolution of this last issue has become even more satisfactory in the very recent analysis of [45].

The aim of the present work is to review this exciting path which characterized the black hole entropy calculation in LQG, trying to show how the analysis of isolated horizons in classical GR, the theory of quantum geometry, and the Chern-Simons theory fuse together to provide a coherent description of the quantum states of isolated horizons, accounting for the entropy. We will present both the  $U(1)$  symmetry reduced and the fully  $SU(2)$  invariant approaches, showing their common features but also their different implications in the quantum theory.

We start by reviewing in Section 2 the formal definition of isolated horizon, through the introduction of the notion of non-expanding horizon first and weakly isolated horizon afterwards. In the second part of this Section, we also provide a classification of IH according to their symmetry group and a notion of staticity condition. We end Section 2 by stating the main equations implied by the isolated horizon boundary conditions for fields at

the horizon, both in the spherically symmetric and the distorted cases.

In Section 3 we construct the conserved symplectic structure of gravity in the presence of a static generic IH. We first use the vector-like (Palatini) variables and then introduce the real (Ashtekar-Barbero) connection variables, showing how, in this passage, a Chern-Simons boundary term appears in the conserved symplectic structure. In Section 4 we briefly review the derivation of the zeroth and first law of isolated horizons.

In Section 5 we show how the classical Hamiltonian framework together with the quantum theory of geometry provide the two pieces of information needed for quantization of Chern-Simons theory on a punctured surface, which describes the quantum degrees of freedom on the horizon. We first present the  $U(1)$  quantization for spherically symmetric horizons and then the  $SU(2)$  scheme for the generic case of distorted horizons, showing how the spherically symmetric picture can be recovered from it.

In Section 6 we perform the entropy calculation of the quantum system previously defined. We first present the powerful methods that have been developed for the resolution of the counting problem in the infinite Chern-Simons level case, involving the  $U(1)$  classical representation theory; in the second part of the Section, we introduce the finite level counting problem by means of the quantum group  $U_q(su(2))$  representation theory, following less rigorous methods. The main results of and differences between the two approaches are analyzed.

In Section 7 we comment on the nature of the entropy degrees of freedom counted in the previous Section and try to compare different points of view appeared in the literature. Section 8 clarifies the role of the Barbero-Immirzi parameter in the LQG black hole entropy calculation, trying to emphasize how the tuning to a given numerical value is not a necessary step to recover the semi-classical area law.

In Section 9 we want to enlighten the connection between the boundary theory and conformal field theory, motivated by other approaches to the entropy problem. In Section 10 we present some recent results [46] on the possibility of using observable effects derived from the black hole entropy description in LQG to experimentally probe the theory, based on some very recent Monte Carlo simulations. Concluding remarks are presented in Section 11.

## 2 Definition of Isolated Horizons

In this Section, we are going to introduce first the notion of Non-Expanding Horizons (NEH) from which, after the imposition of further boundary conditions, we will be able to define Weakly Isolated Horizons (WIH) and the stronger notion of Isolated Horizons (IH), according to [21, 47, 48, 49, 50, 51]. Despite the imposition of these boundary conditions, all these definitions are

far weaker than the notion of an event horizon: The definition of WIH (and IH) extracts from the definition of Killing horizon just that ‘minimum’ of conditions necessary for analogues of the laws of black hole mechanics to hold. Moreover, boundary conditions refer only to behavior of fields at the horizon and the general spirit is very similar to the way one formulates boundary conditions at null infinity.

In the rest of the paper we will assume all manifolds and fields to be smooth. Let  $\mathcal{M}$  be a 4-manifold equipped with a metric  $g_{ab}$  of signature  $(-, +, +, +)$ . We denote  $\Delta$  a null hypersurface of  $(\mathcal{M}, g_{ab})$  and  $\ell$  a future-directed *null* normal to  $\Delta$ . We define  $q_{ab}$  the degenerate intrinsic metric corresponding to the pull-back of  $g_{ab}$  on  $\Delta$ . Denoted  $\nabla_a$  the derivative operator compatible with  $g_{ab}$ , the expansion  $\theta_{(\ell)}$  of a specific null normal  $\ell$  is given by  $\theta_{(\ell)} = q^{ab}\nabla_a\ell_b$ , where the tensor  $q^{ab}$  on  $\Delta$  is the inverse of the intrinsic metric  $q_{ab}$ . With this structure at hand, we can now introduce the definition of NEH.

**Definition:** The internal null boundary  $\Delta$  of an history  $(\mathcal{M}, g_{ab})$  will be called a *non-expanding horizon* provided the following conditions hold:

- i)  $\Delta$  is topologically  $S^2 \times R$ , foliated by a family of 2-spheres  $H$ ;
- ii) The expansion  $\theta_{(\ell)}$  of  $\ell$  within  $\Delta$  vanishes for any null normal  $\ell$ ;
- iii) All field equations hold at  $\Delta$  and the stress-energy tensor  $T_{ab}$  of matter at  $\Delta$  is such that  $-T^a{}_b\ell^b$  is causal and future directed for any future directed null normal  $\ell$ .

Note that if conditions (ii) and (iii) hold for one null normal  $\ell$  they hold for all. Let us discuss the physical meaning of these conditions. The first and the third conditions are rather weak. In particular, the restriction on topology is geared to the structure of horizons that result from gravitational collapse, while the energy condition is satisfied by all matter fields normally considered in general relativity (since it is implied by the stronger dominant energy condition that is typically used). The main condition is therefore the second one, which implies that the *horizon area* ( $a_H$ ) is constant ‘in time’ without assuming the existence of a Killing field.

In the following it will be useful to introduce a null-tetrad which can be built from the null normal field  $\ell^a$  by adding a complex null vector field  $m^a$  tangential to  $\Delta$  and a real, future directed null field  $n^a$  transverse to  $\Delta$  so that the following relations hold:  $n \cdot \ell = -1, m \cdot \bar{m} = 1$  and all other scalar products vanish. The quadruplet  $(\ell, n, m, \bar{m})$  constitutes a null-tetrad. There is, of course, an infinite number of null tetrads compatible with a given  $\ell$ , related to one another by restricted Lorentz rotations. All the conclusions of this Section will not be sensitive to this gauge-freedom.




Figure 1: The characteristic data for a (vacuum) spherically symmetric isolated horizon corresponds to Reissner-Nordstrom data on  $\Delta$ , and free radiation data on the transversal null surface with suitable fall-off conditions. For each mass, charge, and radiation data in the transverse null surface there is a unique solution of Einstein-Maxwell equations locally in a portion of the past domain of dependence of the null surfaces. This defines the phase-space of Type I isolated horizons in Einstein-Maxwell theory. The picture shows two Cauchy surfaces  $M_1$  and  $M_2$  “meeting” at space-like infinity  $i_0$ . A portion of  $\mathcal{I}^+$  and  $\mathcal{I}^-$  are shown; however, no reference to future time-like infinity  $i^+$  is made as the isolated horizon need not to coincide with the black hole event horizon.

Conditions (i) and (iii) also imply that the null normal field  $\ell^a$  is geodesic, i.e., denoting the acceleration of  $\ell^a$  by  $\kappa_{(\ell)}$ , it holds

$$\ell^b \nabla_b \ell^a = \kappa_{(\ell)} \ell^a. \quad (3)$$

The function  $\kappa_{(\ell)}$  is called the *surface gravity* and is not a property of the horizon  $\Delta$  itself, but of a specific null normal to it: if we replace  $\ell$  by  $\ell' = f\ell$ , then the surface gravity becomes  $\kappa_{(\ell')} = f\kappa_{(\ell)} + \mathcal{L}_\ell f$ , where  $\mathcal{L}$  indicates the Lie derivative.

As we saw above, condition (ii) that  $\ell^a$  be expansion-free is equivalent to asking that the area 2-form of the 2-sphere cross-sections of  $\Delta$  be constant in time. This, combined with the Raychaudhuri equation and the matter condition (iii), implies that  $\ell^a$  is also shear-free, namely  $\sigma = 0$ , where  $\sigma = m^a m^b \nabla_a \ell_b$  is the shear of  $\ell$  in the given null tetrad. This in turn implies that the Levi-Civita derivative operator  $\nabla$  compatible with  $g_{ab}$  naturally determines a derivative operator  $D_a$  intrinsic to  $\Delta$  via  $X^a D_a Y^b \equiv X^a \nabla_a Y^b$ , where  $X^a, Y^a \in T(\Delta)$  are tangent to  $\Delta$ . However, since the induced metric  $q_{ab}$  on  $\Delta$  is degenerate, there exist infinitely many derivative operators compatible with it. In order to show that every NEH has a unique intrinsic derivative operator  $D$ , we observe that there is a natural connection 1-form on  $\Delta$ : Since  $\ell$  is expansion, shear and twist free, there exists a one-form  $\omega_a$  intrinsic to  $\Delta$  such that

$$D_a \ell^b = \omega_a \ell^b. \quad (4)$$




Figure 2: Space-times with isolated horizons can be constructed by solving the characteristic initial value problem on two intersecting null surfaces,  $\Delta$  and  $\mathcal{N}$  which intersect in a 2-sphere  $H$ . The final solution admits  $\Delta$  as an isolated horizon [52]. Generically, there is radiation arbitrarily close to  $\Delta$  and no Killing fields in any neighborhood of  $\Delta$ . Note that  $\Psi_4$  need not vanish in any region of space-time, not even on  $\Delta$ .

which in turn implies, for the pull-back on  $\Delta$ ,

$$D_a \ell_b = 0. \quad (5)$$

Relation (5) has two important consequences. Firstly, it is exactly the condition that guarantees that every NEH has a unique intrinsic derivative operator  $D$  [51]. Secondly, it implies that the entire pull-back  $q_{ab}$  of the metric to the horizon is Lie dragged by  $\ell^a$ , namely

$$\mathcal{L}_\ell q_{ab} = 0. \quad (6)$$

From eq. (4) it is immediate to see that the surface gravity  $\kappa_{(\ell)}$  can be written as

$$\kappa_{(\ell)} = \omega_a \ell^a. \quad (7)$$

In terms of the Weyl tensor components, using the Newman-Penrose notation, the boundary conditions (i)-(iii) together with (4) imply that on  $\Delta$  [48]

$$\Psi_0 = C_{abcd} \ell^a m^b \ell^c m^d = 0 \quad \Psi_1 = C_{abcd} \ell^a m^b \ell^c n^d = 0, \quad (8)$$

and hence  $\Psi_2$  is gauge invariant on  $\Delta$ , i.e. independent of the choice of the null-tetrad vectors  $(n, m, \bar{m})$ . The  $\Psi_2$  component of the Weyl curvature will play an important role in the following, entering the expression of some constraints to be satisfied by fields at the horizon (Section 2.3); moreover, its imaginary part encodes the gravitational contribution to the angular-momentum at  $\Delta$  [50] and this will provide a condition for classification of isolated horizons (Section 2.2). A useful relation valid on  $\Delta$  between the intrinsic derivative operator  $D$  and this component of the Weyl tensor is expressed by the exterior derivative of the connection  $\omega$  (which is independent of the choice of  $\ell$ , even if the connection itself is), namely [48]

$$d\omega = 2\text{Im}(\Psi_2)^2 \epsilon, \quad (9)$$

where  ${}^2\epsilon \equiv im \wedge \bar{m}$  is a natural area 2-form on  $\Delta$  ( ${}^2\epsilon$  can be invariantly defined [48]).

So far, we have seen that, even though the three boundary conditions in the definition of NEH are significantly weaker than requiring the horizon to be a Killing horizon for a local Killing vector field, they are strong enough to prove that every null normal vector  $\ell$  is an infinitesimal symmetry for the intrinsic metric  $q$ . It is important though to stress that, on the other hand, space-time  $g_{ab}$  need not admit any Killing field in any neighborhood of  $\Delta$ ; boundary conditions (i)-(iii) refer only to behavior of fields at  $\Delta$ . Note that, at this stage, the only geometric structure intrinsic to  $\Delta$  which is ‘time-independent’ is the metric  $q$ , but not the derivative operator  $D$ . Moreover, since  $\ell$  can be rescaled by a positive definite function, the surface gravity  $\kappa_{(\ell)}$  does not need to be constant on  $\Delta$ . Therefore, additional restriction on the fields at  $\Delta$  need to be introduced in order to establish the *0th* law of black hole mechanics. This will lead us in a moment to the definition of WIH.

The next natural step to strengthen the boundary conditions and restrict the choice of  $\ell$  is to add to the geometric structures conserved along  $\Delta$  also the ‘extrinsic curvature’, once an appropriate definition of it is introduced (since we are dealing with a null surface). In order to do this, we are now going to introduce the definition of WIH and then show how, with this definition, the invariance of a tensor field, which can be thought of as the analogue of the extrinsic curvature, under the infinitesimal transformations generated by a preferred equivalence class  $[\ell]$  can be proven.

**Definition:** A *weakly isolated horizon*  $(\Delta, [\ell])$  consists of a non-expanding horizon  $\Delta$ , equipped with an equivalence class  $[\ell]$  of null normals to it satisfying

- iv)  $\mathcal{L}_\ell \omega = 0$  for all  $\ell \in [\ell]$ , where two future-directed null normals  $\ell$  and  $\ell'$  belong to the same equivalence class  $[\ell]$  if and only if  $\ell' = c\ell$  for some positive constant  $c$ .

Note that, under this constant rescaling, the connection 1-form  $\omega$  is unchanged ( $\omega' = \omega$ <sup>2</sup>) and, therefore, if condition (iv) holds for one  $\ell$ , it holds for all  $\ell$  in  $[\ell]$ . Even though we don’t have a single  $\ell$  yet, by definition, a WIH is equipped with a specific equivalence class  $[\ell]$  of null normals. In particular, given any NEH  $\Delta$ , one can always select an equivalence class  $[\ell]$  of null normals such that  $(\Delta, [\ell])$  is a WIH.

WIH admit a natural, generically unique foliation which can be regarded as providing a ‘rest frame’ for the horizon. As shown in [51], this foliation

---

<sup>2</sup>Under the rescaling  $\ell' \rightarrow f\ell$  the connection 1-form  $\omega$  transforms according to  $\omega_a \rightarrow \omega'_a = \omega_a + D_a \ln f$ .

into *good cuts*<sup>3</sup> always exists and is invariantly defined in the sense that it can be constructed entirely from structures naturally available on  $(\Delta, [\ell])$ . In particular, if the space-time admits an isometry which preserves the given WIH, good cuts are necessarily mapped in to each other by that isometry. We require that the fixed foliation of the horizon coincide with a foliation into ‘good cuts’.

Before showing how, with this further restriction, the *0th* law can now be recovered, let us shortly discuss the physical interpretation of condition (iv). Recall that, on a space-like hypersurface  $H$ , the extrinsic curvature can be defined on  $H$  as  $K_a{}^b = \nabla_a n^b$ , where  $n$  is the unit normal. A natural analog of the extrinsic curvature on a WIH is then  $L_a{}^b = D_a \ell^b$ . By virtue of (4) then, condition (iv) is enough to show that  $L_a{}^b$  is Lie-dragged along  $\ell$ , in fact  $\mathcal{L}_\ell K_a{}^b = \mathcal{L}_\ell(\omega_a \ell^b) = (\mathcal{L}_\ell \omega_a) \ell^b = 0$ . Thus, on a WIH not only the intrinsic metric  $q$  is ‘time-independent’, but also the analog of extrinsic curvature. Note however that the full connection  $D$  or curvature components such as  $\Psi_4$  can be time-dependent on a WIH (see Figure 2).

We are now ready to show that the boundary conditions entering the definition of WIH are enough to prove that the surface gravity is constant on  $\Delta$ , i.e. the *0th* law holds for weakly isolated horizons. By construction, it is immediate to see that  $\ell \cdot {}^2\epsilon = 0$ , and this, together with (9), implies

$$\ell \cdot d\omega = 0; \tag{10}$$

on the other hand,

$$0 = \mathcal{L}_\ell \omega = d(\ell \cdot \omega) + \ell \cdot d\omega. \tag{11}$$

Therefore, by virtue of (7), we have

$$d(\ell \cdot \omega) = d\kappa_{(\ell)} = 0. \tag{12}$$

Thus, surface gravity is constant on  $\Delta$  without requiring the presence of a Killing field even in a neighborhood of  $\Delta$ .

As observed above, in the passage from NEH to WIH we had to introduce a more rigid structure in order to recover the *0th* law of black holes mechanics: whereas on a NEH we only ask that the null normal be a Killing field for the intrinsic metric  $q_{ab}$  on  $\Delta$ , on a WIH, the permissible null normals Lie drag also the connection 1-form  $\omega$ , constraining only certain components of the derivative operator  $D$  to be ‘time-independent’. To see this, we notice that the boundary condition (iv) can be reformulated as

$$[\mathcal{L}_\ell, D]\ell^a = 0 \quad \text{on } \Delta. \tag{13}$$

It is immediate to see that the previous condition implies  $\mathcal{L}_\ell \omega = 0$  through (4). A stronger notion of isolation can now be introduced by requiring

---

<sup>3</sup>A 2-sphere cross-section  $H$  of  $\Delta$  is called a ‘good cut’ if the pull-back of  $\omega_a$  to  $H$  is divergence free with respect to the pull-back of  $g_{ab}$  to  $H$ .

the intrinsic metric  $q$  and the full derivative operator  $D$  (rather than just the 1-form  $\omega$ ) be conserved along  $\Delta$ . This can be achieved by relaxing the restriction of the action of the left side of (13) to  $\ell$  and leads to the definition of isolated horizons.

**Definition:** An *isolated horizon* is a pair  $(\Delta, [\ell])$ , where  $\Delta$  is a NEH equipped with an equivalence class  $[\ell]$  of null normals such that

$$\text{v) } [\mathcal{L}_\ell, D_b]v^a = 0, \text{ for all vector fields } v^a \text{ tangential to } \Delta \text{ and all } \ell \in [\ell].$$

If this condition holds for one  $\ell$  it holds for all  $\ell \in [\ell]$ . Let  $\Delta$  be a NEH with geometry  $(q, D)$ . We will say that this geometry admits an *isolated horizon structure* if there exists a null normal  $\ell$  satisfying (v). Intuitively, a NEH is an IH if the entire geometry  $(q, D)$  of the NEH is ‘time-independent’. From the perspective of the intrinsic geometry, this is a stronger and perhaps more natural notion of ‘isolation’ than that captured in the definition of a WIH. However, unlike (iv), condition (v) is a genuine restriction. In fact, while any NEH can be made a WIH simply by choosing an appropriate class  $\ell$  of null normals, not every NEH admits a null normal satisfying (v) and generically this condition does suffice to single out the equivalence class  $[\ell]$  uniquely [51]. Thus, even though (v) is a stronger condition than (iv), it is still *very* weak compared to conditions normally imposed: using the initial value problem based on two null surfaces [53], it can be shown that the definition of IH contains an infinite-dimensional class of other examples [52]. In particular, while all geometric fields on  $\Delta$  are time-independent as on a Killing horizon, the field  $\Psi_4$ , for example, can be ‘time-dependent’ on a generic IH.

To summarize, IH are null surfaces, foliated by a (preferred) family of marginally trapped 2-spheres such that certain geometric structures intrinsic to  $\Delta$  are time independent. The presence of trapped surfaces motivates the term ‘horizon’ while the fact that they are *marginally* trapped — i.e., that the expansion of  $\ell^a$  vanishes — accounts for the adjective ‘isolated’. The definition extracts from the definition of Killing horizon just that ‘minimum’ of conditions necessary for analogues of the laws of black hole mechanics to hold<sup>4</sup>.

**Remarks:**

1. All the boundary conditions are satisfied by stationary black holes in the Einstein-Maxwell-dilaton theory possibly with cosmological constant. More importantly, starting with the standard stationary black

---

<sup>4</sup>We will see in the following how the first law can be recovered by requiring the time evolution along vector fields  $t^a \in T(\mathcal{M})$ , which are time translations at infinity and proportional to the null generators  $\ell$  at the horizon, to correspond to a Hamiltonian time evolution [48]

holes, and using known existence theorems one can specify procedures to construct new solutions to field equations which admit isolated horizons as well as radiation at null infinity [52]. These examples already show that, while the standard stationary solutions have only a finite parameter freedom, the space of solutions admitting IH is *infinite* dimensional. Thus, in the Hamiltonian picture, even the reduced phase-space is infinite dimensional; the conditions thus admit a very large class of examples. Nevertheless, space-times admitting IH are very special among generic members of the full phase-space of general relativity. The reason is apparent in the context of the characteristic formulation of general relativity where initial data are given on a set (pairs) of null surfaces with non trivial domain of dependence. Let us take an isolated horizon as one of the surfaces together with a transversal null surface according to the diagram shown in Figure 1. Even when the data on the IH may be infinite dimensional, in all cases no transversing radiation data is allowed by the IH boundary condition.

2. The freedom in the choice of the null normal  $\ell$  we saw existing for isolated horizons is present also in the case of Killing horizons. Given a Killing horizon  $\Delta_K$ , surface gravity is defined as the acceleration of a static particle near the horizon, moving on an orbit of a Killing field  $\eta$  normal to  $\Delta_K$ , as measured at spatial infinity. However, if  $\Delta_K$  is a Killing horizon for  $\eta$ , it is also for  $c\eta, \forall c = \text{const} > 0$ . Therefore, surface gravity is not an intrinsic property of  $\Delta_K$ , but depends also on a specific choice of  $\eta$ : its normalization is undetermined, since it scales under constant scalings of the Killing vector  $\eta$  (even though this freedom does not affect the *0th* law). However, even if one cannot normalize  $\eta$  at the horizon (since  $\eta^2 = 0$  there), in the case of asymptotically flat space-times admitting global Killing fields, its normalization can be specified in terms of the behavior of  $\eta$  at infinity. For instance, in the static case, the Killing field  $\eta$  can be canonically normalized by requiring that it have magnitude-squared equal to  $-1$  at infinity. In absence of a global Killing field or asymptotic flatness though, this strategy does not work and one has to keep this constant rescaling freedom in the definition of surface gravity. In the context of isolated horizons, then, it is natural to keep this freedom. Nevertheless, one can, if necessary, select a specific  $\ell$  in  $[\ell]$  by requiring, for instance,  $\kappa_{(\ell)}$  to coincide with the surface gravity of black holes in the Reissner-Nordstrom family:

$$\kappa_{(\ell)} = \frac{\sqrt{(M^2 - Q^2)}}{2M[M + \sqrt{(M^2 - Q^2)}] - Q^2}, \quad (14)$$

where  $M$  is the mass and  $Q$  the electric charge of the black hole. Indeed this choice is the one that makes the zero, and first law of IH look just

as the corresponding laws of stationary black hole mechanics [48, 41] (see Section 4 for a discussion on the zeroth and first laws).

3. Notice that the above definition is completely geometrical and does not make reference to the tetrad formulation. There is no reference to any internal gauge symmetry. In what follows we will deal with general relativity in the first order formulation which will introduce, by the choice of variables, an internal gauge group corresponding to local  $SL(2, \mathbb{C})$  transformations (in the case of Ashtekar variables) or  $SU(2)$  transformations (in the case of real Ashtekar-Barbero variables). As pointed out in the introduction, the original quantization scheme of [19, 49, 54] uses a gauge symmetry reduced framework while a more recent analysis [40, 41, 42] preserves the full internal gauge symmetry. Both approaches are the subject of Sections 5.1 and 5.2.

## 2.1 IH Classification According to Their Symmetry Groups

Next, let us examine symmetry groups of isolated horizons. As seen above, boundary conditions impose restrictions on dynamical fields and also on gauge transformations on the boundary. At infinity all transformations are required to preserve asymptotic flatness; hence, the asymptotic symmetry group reduces to the Poincaré group. On the other hand, a *symmetry* of  $(\Delta, q, D, [\ell^a])$  is a diffeomorphism on  $\Delta$  which preserves the horizon geometry  $(q, D)$  and at most rescales elements of  $[\ell^a]$  by a positive constant. These diffeomorphisms must be compositions of translations along the integral curves of  $\ell^a$  and general diffeomorphisms on a 2-sphere in the foliation. Thus, the boundary conditions reduce the symmetry group  $G_\Delta$  to a semi-direct product of diffeomorphisms generated by  $\ell^a$  with  $\text{Diff}(S^2)$ . In fact, there are only three possibilities for  $G_\Delta$  [50]:

- (a) Type I: the isolated horizon geometry is spherical; in this case,  $G_\Delta$  is four dimensional ( $SO(3)$  rotations plus rescaling-translations<sup>5</sup> along  $\ell$ );
- (b) Type II: the isolated horizon geometry is axisymmetric; in this case,  $G_\Delta$  is two dimensional (rotations round symmetry axis plus rescaling-translations along  $\ell$ );
- (c) Type III: the diffeomorphisms generated by  $\ell^a$  are the only symmetries;  $G_\Delta$  is one dimensional.

Note that these symmetries refer only to the horizon geometry. The full space-time metric need not admit any isometries even in a neighborhood of the horizon.

---

<sup>5</sup>In a coordinate system where  $\ell^a = (\partial/\partial v)^a$  the rescaling-translation corresponds to the affine map  $v \rightarrow cv + b$  with  $c, b \in \mathbb{R}$  constants.

## 2.2 IH Classification According to the Reality of $\Psi_2$

As observed above, the gravitational contribution to angular momentum of the horizon is coded in the imaginary part of  $\Psi_2$  [50]. Therefore, the reality of  $\Psi_2$  allows us to introduce an important classification of isolated horizons.

- (1) *Static*: In the Newman-Penrose formalism (in the null tetrads adapted to the IH geometry introduced above), static isolated horizons are characterized by the condition

$$\text{Im}(\Psi_2) = 0 \tag{15}$$

on the Weyl tensor component  $\Psi_2 = C_{abcd}\ell^a m^b \bar{m}^c n^d$ . This corresponds to having the horizon locally “at rest”. In the axisymmetric case, according to the definition of multiple moments of Type II horizons constructed in [54], static isolated horizons are *non-rotating* isolated horizons, *i.e.* those for which all angular momentum multiple moments vanish. Static black holes (e.g., those in the Reissner-Nordstrom family) have static isolated horizons. There are Type I, II and III static isolated horizons.

- (2) *Non-Static*: In the Newman-Penrose formalism, non-static isolated horizons are characterized by the condition

$$\text{Im}(\Psi_2) \neq 0. \tag{16}$$

The horizon is locally “in motion”. The Kerr black hole is an example of this type.

### Remark:

In the rest of the paper we will concentrate only on static isolated horizons. We will show that for this class of IH one can construct a conserved pre-symplectic structure with no need to make any symmetry assumptions on the horizon. On the other hand, the usual pre-symplectic structure is not conserved in the presence of a non-static black hole (see Section 3.2 below), which implies that a complete treatment of non-static isolated horizons (including rotating isolated horizons) remains open—see [42] for a proposal leading to a conserved symplectic structure for non-static isolated horizons and the restoration of diffeomorphisms invariance.

## 2.3 The Horizon Constraints

We are now going to use the definition of isolated horizons provided above to derive some equations which will play a central role in the sequel. General relativity in the first order formalism is described in terms a tetrad of four

1-forms  $e^I$  ( $I = 0, 3$  internal indices) and a Lorentz connection  $\omega^{IJ} = -\omega^{JI}$ . The metric can be recovered by

$$g_{ab} = e_a^I e_b^J \eta_{IJ} \quad (17)$$

where  $\eta_{IJ} = \text{diag}(-1, 1, 1, 1)$ . In the time gauge, where the tetrad  $e^I$  is such that  $e^0$  is a time-like vector field normal to the Cauchy surface  $M$ , the three 1-forms  $K^i = \omega^{0i}$  play a special role in the parametrization of the phase-space. In particular the so-called Ashtekar connection is

$$A_a^{+i} = \Gamma_a^i + iK_a^i \quad (18)$$

where  $\Gamma^i = -\frac{1}{2}\epsilon^{ijk}\omega_{jk}$  is the spin connection satisfying Cartan's first equation

$$d_\Gamma e^i = 0. \quad (19)$$

On  $\Delta$  one can, of course, express the tetrad  $e^I$  in terms of the null-tetrad  $(\ell, n, m, \bar{m})$  introduced above; in particular, at  $H = \Delta \cap M$ , the normal to  $M$  can be written as  $e^{0a} = (\ell^a + n^a)/\sqrt{2}$  at  $H$  (recall that  $n^a$  is normalized according to  $n \cdot \ell = -1$ ).

We also introduce the 2-form

$$\Sigma^{IJ} \equiv e^I \wedge e^J \quad \text{and} \quad \Sigma^{+i} \equiv \epsilon^i{}_{jk} \Sigma^{jk} + 2i\Sigma^{0i}$$

and  $F^i(A^+)$  the curvature of the connection  $A^{+i}$ . In Section 5.2, at  $H$ , we will also often work in the gauge where  $e^1$  is normal to  $H$  and  $e^2$  and  $e^3$  are tangent to  $H$ . This choice is only made for convenience, as the equations used there are all gauge covariant, their validity in one frame implies their validity in all frames.

When written in connection variables, the isolated horizons boundary condition implies the following relationship between the curvature of the Ashtekar connection  $A^{+i}$  at the horizon and the 2-form  $\Sigma^i \equiv \text{Re}[\Sigma^{+i}] = \epsilon^i{}_{jk} e^j \wedge e^k$  [18, 42]

$$\underline{\underline{F}}_{ab}{}^i(A^+) = (\Psi_2 - \Phi_{11} - \frac{R}{24}) \underline{\underline{\Sigma}}_{ab}^i, \quad (20)$$

where the double arrows denote the pull-back to  $H$ . For simplicity, here we will assume that no matter is present at the horizon,  $\Phi_{11} = R = 0$ , hence

$$\underline{\underline{F}}_{ab}{}^i(A^+) = \Psi_2 \underline{\underline{\Sigma}}_{ab}^i. \quad (21)$$

An important point here is that the previous expression is valid for any two sphere  $S^2$  (not necessarily a horizon) embedded in space-time in an adapted null tetrad where  $\ell^a$  and  $n^a$  are normal to  $S^2$ . In the special case of pure gravity, and due to the vanishing of both the expansion and shear of the

generators congruence  $\ell^a$ , the Weyl component  $\Psi_2$  at the Horizon is simply related to the Gauss scalar curvature  $R^{(2)}$  of the two spheres.

Eq. (20) can be derived starting from the identity (that can be derived from Cartan's second structure equations)

$$F_{ab}{}^i(A^+) = -\frac{1}{4}R_{ab}{}^{cd}\Sigma_{cd}{}^i, \quad (22)$$

where  $R_{abcd}$  is the Riemann tensor, using the null-tetrad formalism (see for instance [55]) with the null-tetrad introduced above, and the definitions  $\Psi_2 = C_{abcd}\ell^am^b\bar{m}^cn^d$  and  $\Phi_{11} = R_{ab}(\ell^an^b + m^a\bar{m}^b)/4$ , where  $R_{ab}$  is the Ricci tensor and  $C_{abcd}$  the Weyl tensor (for an explicit derivation using the spinors formalism see [18] Appendix B).

In the case of Type I IH, we have [18]

$$\Psi_2 - \Phi_{11} - \frac{R}{24} = -\frac{2\pi}{a_H} \quad (23)$$

at  $H$ , where  $a_H$  is the area of the IH. Therefore, the horizon constraint (21) becomes

$$F_{\leftarrow ab}{}^i(A^+) = -\frac{2\pi}{a_H}\Sigma_{\leftarrow ab}{}^i \quad (24)$$

in the spherically symmetric case [18, 41].

Notice that the imaginary part of the eq. (21) implies that, for static IH,

$$d_{\leftarrow\Gamma}K^i = 0. \quad (25)$$

Relation (21), which follows from the boundary conditions on  $\Delta$ , provides a restriction on the possible histories of the phase-space whose points are represented by values of the space-time fields  $(A^+, \Sigma^+)$ . In particular, at  $H$ , the behavior of the (curvature of) Ashtekar connection  $A^+$  is related to the pull-back of  $\Sigma$  through the Weyl tensor component  $\Psi_2$ . In this sense, at the classical level, all the horizon degrees of freedom are encoded in the range of possible values of  $\Psi_2$ , which, without symmetry restriction, can be infinite-dimensional. We will see in the next Sections how this picture changes at the quantum level.

Another important relation, valid for static IH and following from condition (v) [42], is

$$K_{\leftarrow}^j \wedge K_{\leftarrow}^k \epsilon_{ijk} = c \Sigma_{\leftarrow}^i \quad (26)$$

for  $c : H \rightarrow \mathbb{R}$  an extrinsic curvature scalar.

**Remark:**

In the GHP formalism [56], a null tetrad formalism compatible with the IH system, the scalar curvature of the two-spheres normal to  $\ell^a$  and  $n^a$  is given by

$$R^{(2)} = K + \bar{K} \quad (27)$$

where  $K = \sigma\sigma' - \rho\rho' - \Psi_2 + R + \Phi_{11}$ , while  $\sigma$ ,  $\rho$ ,  $\sigma'$  and  $\rho'$  denote spin, shear and expansion spin coefficients associated with  $\ell^a$  and  $n^a$  respectively. The shear-free and expansion-free conditions in the definition of IH translate into  $\rho = 0 = \sigma$  in the GHP formalism, namely

$$R^{(2)} = -2\Psi_2 \quad (28)$$

Similarly, the curvature scalar  $c$  in (26) can be expressed in terms of spin coefficients as

$$c = \frac{1}{2}(\rho'\bar{\rho}' - \sigma'\bar{\sigma}') \quad (29)$$

which is invariant under null tetrad transformations fixing  $\ell^a$  and  $n^a$ .

### 3 The Conserved Symplectic Structure

In this Section we prove the conservation of the symplectic structure of gravity in the presence of an isolated horizon that is not necessarily spherically symmetric but static. For the non-static case, we will see how diffeomorphisms tangent to the horizon are no longer degenerate directions of the symplectic structure and, therefore, the quantization techniques described in Section 5 need to be generalized. Quantization of *rotating* black holes remains an open issue in the framework of LQG.

Conservation of the symplectic structure was first shown in [18] for Type I IH in the  $U(1)$  gauged fixed formalism. In the rest of this Section, we will follow the proof presented in [42], where the full  $SU(2)$  invariant formalism is applied to generic distorted IH. The gauge fixed symplectic form for Type I IH will be introduced at the end of Section 3.3.1 in order to describe the  $U(1)$  quantization of spherically symmetric horizons in Section 5.1.

#### 3.1 Action Principle and Phase-Space

The action principle of general relativity in self dual variables containing an inner boundary satisfying the IH boundary condition (for asymptotically flat space-times) takes the form

$$S[e, A^+] = -\frac{i}{\kappa} \int_{\mathcal{M}} \Sigma_i^+(e) \wedge F^i(A^+) + \frac{i}{\kappa} \int_{\tau_\infty} \Sigma_i^+(e) \wedge A^{+i}, \quad (30)$$

where  $\kappa = 16\pi G$  and a boundary contribution at a suitable time cylinder  $\tau_\infty$  at infinity is required for the differentiability of the action. No horizon boundary term is necessary if one allows variations that fix an isolated horizon geometry up to diffeomorphisms and Lorentz transformations. This is a very general property, as shown in [41].

First variation of the action yields

$$\begin{aligned}\delta S[e, A^+] &= \frac{-i}{\kappa} \int_{\mathcal{M}} \delta \Sigma_i^+(e) \wedge F^i(A^+) - d_{A^+} \Sigma_i^+ \wedge \delta A^{+i} + d(\Sigma_i^+ \wedge \delta A^{+i}) \\ &+ \frac{i}{\kappa} \int_{\tau_\infty} \delta(\Sigma_i^+(e) \wedge A^{+i}),\end{aligned}\quad (31)$$

from which the self dual version of Einstein's equations follow

$$\begin{aligned}\epsilon_{ijk} e^j \wedge F^i(A^+) + i e^0 \wedge F_k(A^+) &= 0 \\ e_i \wedge F^i(A^+) &= 0 \\ d_{A^+} \Sigma_i^+ &= 0\end{aligned}\quad (32)$$

as the boundary terms in the variation of the action cancel.

We denote  $\Gamma$  the phase-space of a space-time manifold with an internal boundary satisfying the boundary condition corresponding to static IH, and asymptotic flatness at infinity. The phase-space of such system is defined by an infinite dimensional manifold where points  $p \in \Gamma$  are given by solutions to Einstein's equations satisfying the static IH boundary conditions. Explicitly, a point  $p \in \Gamma$  can be parametrized by a pair  $p = (\Sigma^+, A^+)$  satisfying the field equations (32) and the requirements of the IH definition provided above. In particular fields at the boundary satisfy Einstein's equations and the constraints given in Section 2.3. Let  $T_p(\Gamma)$  denote the space of variations  $\delta = (\delta \Sigma^+, \delta A^+)$  at  $p$  (in symbols  $\delta \in T_p(\Gamma)$ ). A very important point is that the IH boundary conditions severely restrict the form of field variations at the horizon. Thus we have that variations  $\delta = (\delta \Sigma^+, \delta A^+) \in T_p(\Gamma)$  are such that for the pull-back of fields on the horizon they correspond to linear combinations of  $SL(2, \mathbb{C})$  internal gauge transformations and diffeomorphisms preserving the preferred foliation of  $\Delta$ . In equations, for  $\alpha : \Delta \rightarrow sl(2, \mathbb{C})$  and  $v : \Delta \rightarrow T(H)$  we have that

$$\begin{aligned}\delta \Sigma^+ &= \delta_\alpha \Sigma^+ + \delta_v \Sigma^+ \\ \delta A^+ &= \delta_\alpha A^+ + \delta_v A^+\end{aligned}\quad (33)$$

where the infinitesimal  $SL(2, \mathbb{C})$  transformations are explicitly given by

$$\delta_\alpha \Sigma^+ = [\alpha, \Sigma^+], \quad \delta_\alpha A^+ = -d_{A^+} \alpha, \quad (34)$$

while the diffeomorphisms tangent to  $H$  take the following form

$$\begin{aligned}\delta_v \Sigma_i^+ &= \mathcal{L}_v \Sigma_i^+ = \underbrace{v \lrcorner d_{A^+} \Sigma_i^+}_{= 0 \text{ (Gauss)}} + d_{A^+} (v \lrcorner \Sigma^+)_i - [v \lrcorner A^+, \Sigma^+]_i \\ \delta_v A^{+i} &= \mathcal{L}_v A^{+i} = v \lrcorner F^{+i} + d_{A^+} (v \lrcorner A^+)^i,\end{aligned}\quad (35)$$

where  $(v \lrcorner \omega)_{b_1 \dots b_{p-1}} \equiv v^a \omega_{ab_1 \dots b_{p-1}}$  for any  $p$ -form  $\omega_{b_1 \dots b_p}$ , and the first term in the expression of the Lie derivative of  $\Sigma_i^+$  can be dropped due to the Gauss constraint  $d_A \Sigma_i^+ = 0$ .

### 3.2 The Conserved Symplectic Structure in Terms of Vector Variables

So far we have defined the covariant phase-space as an infinite dimensional manifold. For it to become a phase-space it is necessary to provide it with a pre-symplectic structure. As the field equations, the pre-symplectic structure can be obtained from the first variation of the action (32). In particular a symplectic potential density for gravity can be directly read off from the total differential term in (32) [57, 58]. In terms of the Ashtekar connection and the densitized tetrad, the symplectic potential density is

$$\theta(\delta) = \frac{-i}{\kappa} \delta_{[1} \Sigma_i^+ \wedge \delta A^{+i} \quad \forall \delta \in T_p \Gamma \quad (36)$$

and the symplectic current takes the form

$$J(\delta_1, \delta_2) = -\frac{2i}{\kappa} \delta_{[1} \Sigma_i^+ \wedge \delta_2] A^{+i} \quad \forall \delta_1, \delta_2 \in T_p \Gamma. \quad (37)$$

Einstein's equations imply  $dJ = 0$ . From Stokes theorem applied to the four dimensional (shaded) region in Figure 1 bounded by  $M_1$  in the past,  $M_2$  in the future, a time-like cylinder at spacial infinity on the right, and the isolated horizon  $\Delta$  on the left, it can be shown that the symplectic form

$$\kappa \Omega_M(\delta_1, \delta_2) = \int_M \delta_{[1} \Sigma^i \wedge \delta_2] K_i \quad (38)$$

is conserved in the sense that  $\Omega_{M_2}(\delta_1, \delta_2) = \Omega_{M_1}(\delta_1, \delta_2)$ , where  $M$  is a Cauchy surface representing space. The symplectic form above, written in terms of the vector-like (or Palatini) variables  $(\Sigma, K)$ , is manifestly real and has no boundary contribution.

In the case of diffeomorphisms for the variations on the horizon, conservation of the symplectic form (38) follows from the relation

$$v_{\perp} \Sigma_i \wedge K^i = 0, \quad (39)$$

which holds *only* for static IH [42]. Therefore, in presence of a non-static IH, the symplectic form (38) for gravity is no longer conserved: rotating isolated horizons boundary conditions break diffeomorphisms invariance<sup>6</sup>.

---

<sup>6</sup>More precisely, the gauge symmetry content of isolated horizon systems is characterized by the degenerate directions of the pre-symplectic structure. As shown in [42], tangent vectors of the phase-space  $\Gamma$ , i.e. variations  $\delta \in T_p \Gamma$  corresponding to diffeomorphisms tangent to the horizon are degenerate directions of  $\Omega_M$  if and only if the isolated horizon is static. Nevertheless, variations corresponding to  $SU(2)$  gauge transformations remain degenerate directions also in the non-static case.

### 3.3 The Conserved Symplectic Structure in Terms of Real Connection Variables

In order to be able to apply the LQG formalism in Section 5 to quantize the bulk theory, we now want to introduce the Ashtekar-Barbero variables

$$A_a^i = \Gamma_a^i + \beta K_a^i \quad (40)$$

where  $\beta$  is the Barbero-Immirzi parameter. We can write the symplectic potential corresponding to (38) as

$$\begin{aligned} \kappa \Theta(\delta) &= \frac{1}{\beta} \int_M \Sigma_i \wedge \delta(\Gamma^i + \beta K^i) - \frac{1}{\beta} \int_M \Sigma_i \wedge \delta \Gamma^i \\ &= \frac{1}{\beta} \int_M \Sigma_i \wedge \delta(\Gamma^i + \beta K^i) + \frac{1}{\beta} \int_H e_i \wedge \delta e^i, \end{aligned} \quad (41)$$

where in the last line we have used a very important property of the spin connection [13] compatible with  $e^i$ , namely

$$\int_M \Sigma_i \wedge \delta \Gamma^i = \int_H -e_i \wedge \delta e^i. \quad (42)$$

In terms of the Ashtekar-Barbero connection the symplectic structure (38) takes the form

$$\kappa \Omega_M(\delta_1, \delta_2) = \frac{1}{\beta} \int_M \delta_{[1} \Sigma^i \wedge \delta_2] A_i - \frac{1}{\beta} \int_H \delta_{[1} e^i \wedge \delta_2] e_i. \quad (43)$$

Before introducing connection variables also for the boundary theory, a some comments are now in order. We have shown that in the presence of a static isolated horizon the conserved pre-symplectic structure is the usual one when written in terms of vector-like variables. When we write the pre-symplectic structure in terms of Ashtekar-Barbero connection variables in the bulk, the pre-symplectic structure acquires a boundary term at the horizon of the simple form [41, 59]

$$\kappa \Omega_H(\delta_1, \delta_2) = \frac{1}{\beta} \int_H \delta_{[1} e^i \wedge \delta_2] e_i. \quad (44)$$

This boundary contribution provides an interesting insight already at the classical level, as the boundary symplectic structure, written in this way, has a remarkable implication for geometric quantities of interest in the first order formulation. More precisely, this implies the kind of non-commutativity of flux variables that is compatible with the use of the holonomy-flux algebra as the starting point for LQG quantization. In fact, (44) implies  $\{e_a^i(x), e_b^j(y)\} = \epsilon_{ab} \delta^{ij} \delta(x, y)$  from which one can compute the Poisson brackets among surface fluxes

$$\Sigma(S, \alpha) = \int_{S \subset H} \text{Tr}[\alpha \Sigma], \quad (45)$$

where  $S \subset H$  and  $\alpha : H \rightarrow \mathfrak{su}(2)$ , and see that they reproduce the  $\mathfrak{su}(2)$  Lie algebra [41]. This is an interesting property that follows entirely from classical considerations using smooth field configurations. This fact strengthens even further the relevance of the uniqueness theorems [60], as they assume the use of the *holonomy-flux* algebra as the starting point for quantization, for which flux variables satisfy commutation relations corresponding exactly to this Poisson structure.

A second observation is that the symplectic term (44) shows that the boundary degrees of freedom could be described in terms of the pull back of the triad fields  $e^i$  on the horizon subjected to the obvious constraint

$$\Sigma_H^i = \Sigma_{Bulk}^i, \quad (46)$$

which are three first class constraints—as it follows from (44)—for the six unconstrained phase-space variables  $e^i$ . One could try to quantize the IH system in this formulation in order to address the question of black hole entropy calculation. Despite the non-immediacy of the background independent quantization of the boundary theory in terms of triad fields, as pointed out in [42], difficulties would appear in the quantum theory due to the presence of degenerate geometry configurations which would constitute residual gauge local degrees of freedom in  $e^i$  not killed by the quantum imposition of (46). This would naively lead to an infinite entropy.

While a more detailed study of the quantization of the  $e^i$  fields on  $H$  would be definitely interesting and might reveal interesting geometric implications, the situation is very much reminiscent of the theory in the bulk, where the same issue of choice of continuum variables to use for the phase-space parametrization appears. More precisely, while the bulk theory can very well be described in terms of vector-like variables  $(\Sigma, K)$ , we wouldn't know how to quantize the theory in a background independent framework using these variables. That is why we chose a phase-space parametrization in terms of (Ashtekar-Barbero) connections and then apply the LQG machinery to quantize the bulk theory. This suggests that also for the boundary theory the passage to connection variables may simplify the quantization process<sup>7</sup>. Evidence for this comes, for instance, from the spherically symmetric case, where the degrees of freedom are encoded instead in a connection  $A^i$  and the analog of the constraints  $\Sigma_H^i = 0$  (where there are no bulk punctures) are  $F^i(A) = 0$  (see the following Subsection for more details). The dimensionality of both the unconstrained phase space and constraint surfaces are the same as in the treatment based on triads; however, the constraint  $F^i(A) = 0$  completely annihilates the local degrees of freedom at places where there are no punctures, rendering the entropy finite. This motivates the use of connection variables to describe (44). In the generic distorted

---

<sup>7</sup>Recall that the boundary theory was originally derived in terms of connection variables [18, 19].

case, this alternative description can be achieved by the introduction of a pair of  $SU(2)$  connection variables

$$A_{\sigma_+}^i = \Gamma^i + \frac{2\pi}{a_H} \sigma_+ e^i \quad \text{and} \quad A_{\sigma_-}^i = \Gamma^i + \frac{2\pi}{a_H} \sigma_- e^i, \quad (47)$$

where  $\sigma_{\pm}$  are two new free adimensional parameters (the factor  $1/a_H$  is there for dimensional reasons). In terms of these new variables, the boundary contribution of the conserved symplectic form (44) becomes

$$\kappa\beta\Omega_H(\delta_1, \delta_2) = \frac{a_H}{2\pi(\sigma_-^2 - \sigma_+^2)} \left( \int_H \delta_{[1} A_{\sigma_+}^i \wedge \delta_{2]} A_{\sigma_+ i} - \int_H \delta_{[1} A_{\sigma_-}^i \wedge \delta_{2]} A_{\sigma_- i} \right).$$

See [42] for the proof of the previous relation. From the IH boundary conditions, through the relations (21) and (26), Cartan's equations, and the definitions (47), the following relations for the new variables hold

$$F_{\leftarrow}^i(A_{\sigma_{\pm}}) = \Psi_2 \Sigma_{\leftarrow}^i + \left( \frac{\pi}{a_H} \sigma_{\pm}^2 + \frac{c}{2} \right) \Sigma_{\leftarrow}^i. \quad (48)$$

This means that there is a two-parameter family of equivalent classical descriptions of the system that in terms of triad variables is described by (44) (we will see in the sequel that the two parameter freedom reduces indeed to a single one when additional consistency requirements are taken into account). The appearance of these new parameters  $\sigma_{\pm}$  is strictly related to the introduction of the  $SU(2)$  connection variables (as was already observed in [41]). This fact fully reflects the analogy with the bulk theory, where the appearance of the Barbero-Immirzi parameter follows from the introduction of the Ashtekar-Barbero variables (replacing the vector-like variables) in the parametrization of the phase-space of general relativity in the bulk.

In the quantum theory, at points where there are no punctures from the bulk, the two connections are subjected to the six first class constraints  $F^i(A_{\gamma}) = 0 = F^i(A_{\sigma})$  implying the absence of local degrees of freedom at these places. The new variables resolve in this way the difficulty related to the treatment in terms of the triads  $e^i$ . In addition, the connection fields  $A_{\gamma}$  and  $A_{\sigma}$  are described by Chern-Simons symplectic structures respectively, with levels

$$k_+ = -k_- = \frac{a_H}{4\pi\ell_p^2\beta(\sigma_-^2 - \sigma_+^2)}, \quad (49)$$

which will allow us the use of some of the standard techniques, firstly applied to Type I isolated horizons in [19], for the quantization of arbitrary static isolated horizons.

### 3.3.1 The Spherically Symmetric Case

Before starting the discussion on the boundary theory quantization, we are now going to revise briefly the description of the horizon degrees of freedom

in terms of connection variables for Type I IH. We will first derive the boundary theory in the full  $SU(2)$  invariant set-up and at the end present its  $U(1)$  gauge fixed formulation.

For a spherically symmetric IH the Weyl tensor component  $\Psi_2$  and the curvature invariant  $c$  in (26) take the constant values:  $\Psi_2 = -\frac{2\pi}{a_H}$  and  $c = \frac{2\pi}{a_H}$  (see [41]), from which the horizon constraint (21) and the relation (26) become

$$F_{\leftarrow ab}^i(A^+) = -\frac{2\pi}{a_H} \Sigma_{\leftarrow ab}^i \quad \text{and} \quad \epsilon^i{}_{jk} K_{\leftarrow}^j \wedge K_{\leftarrow}^k = \frac{2\pi}{a_H} \Sigma_{\leftarrow}^i. \quad (50)$$

The previous equations in turn imply that the curvature of an  $SU(2)$  connection  $A_\sigma^i = \Gamma^i + \sigma K^i$  on the horizon be related to the pull-back of the 2-form  $\Sigma$  in the bulk through

$$F_{\leftarrow}^i(A_\sigma) = -\frac{\pi(1-\sigma^2)}{a_H} \Sigma_{\leftarrow}^i, \quad (51)$$

where a new parameter  $\sigma$  (independent of the Barbero-Immirzi parameter) has appeared due to the introduction of the boundary connection  $A_\sigma^i$ . We can now express the symplectic form boundary contribution (44) in terms of this new connection. If we do so, the symplectic structure of spherically symmetric IH takes the form [41]

$$\kappa\beta \Omega_M(\delta_1, \delta_2) = \int_M \delta_{[1} \Sigma^i \wedge \delta_{2]} A_i - \frac{a_H}{\pi(1-\sigma^2)} \int_H \delta_{[1} A_\sigma^i \wedge \delta_{2]} A_{\sigma i}. \quad (52)$$

Therefore, the degrees of freedom of Type I IH are described by a single  $SU(2)$  Chern-Simons theory with level

$$k = \frac{a_H}{4\pi\ell_p^2\beta(1-\sigma^2)}, \quad (53)$$

which depends on both the Barbero-Immirzi and the new parameter  $\sigma$ .

In [18] the classical description of spherically symmetric IH phase-space was original performed by introducing a partial gauge fixing from the internal gauge  $SL(2, \mathbb{C})$  to  $U(1)$ . In this setting one fixes an internal direction  $r^i \in \mathfrak{su}(2)$  and then the horizon degrees of freedom are encoded on the Abelian part  $W$  of the pull-back to  $H$  of the connection  $A$ , namely

$$W_a \equiv -\frac{1}{\sqrt{2}} \Gamma_{\leftarrow a}^i r_i. \quad (54)$$

The IH boundary condition (51) now becomes

$$dW = -\frac{2\pi}{a_H} \Sigma_{\leftarrow}^i r_i, \quad \Sigma_{\leftarrow}^i x_i = 0, \quad \Sigma_{\leftarrow}^i y_i = 0, \quad (55)$$

where  $x^i, y^i \in \mathfrak{su}(2)$  are arbitrary vectors completing an internal triad; and the horizon conserved symplectic structure takes the form

$$\kappa\beta\Omega_M(\delta_1, \delta_2) = \int_M \delta_{[1}\Sigma^i \wedge \delta_2]A_i - \frac{a_H}{\pi} \int_H \delta_1 W \wedge \delta_2 W. \quad (56)$$

Therefore, in addition to the standard bulk term, the symplectic structure contains a surface term which coincides with that of a  $U(1)$  Chern-Simons theory for  $W$  with level

$$k = \frac{a_H}{4\pi\beta\ell_{\text{P}}^2}. \quad (57)$$

Notice that, in this gauge fixed set-up, no new free parameter appears in the boundary theory description. We will see in the next Section the important role of this additional free parameter introduced by the description of the horizon theory in terms of  $SU(2)$  connections and absent in the  $U(1)$  formulation.

## 4 The Zeroth and First Laws of BH Mechanics for Isolated Horizons

The definition of WIH given in Section 2 implies automatically the zeroth law of BH mechanics as  $\kappa_{(\ell)}$  is constant on  $\Delta$  (see eq. (12)). In turn, the first law cannot be tested unless a definition of energy  $E_H^t$  associated with the IH is given. Since there can be radiation in space-time outside  $\Delta$ ,  $E_{ADM}$  is not a good measure of  $E_H^t$ . In absence of global Killing vector fields, the behavior of the (time) vector fields  $t^a \in T(M)$  near the horizon is unrelated to its behavior near infinity and hence, at the horizon, it is not possible to define a unique time evolution. Fortunately, the Hamiltonian framework provides an elegant way to define a notion of energy associated to the horizon. This consists of requiring the time evolution along vector fields  $t^a$ , which are time translations at infinity and proportional to the null generators  $\ell$  at the horizon, to correspond to a Hamiltonian time evolution [48]. More precisely, denote by  $\delta_t : \Gamma \rightarrow T(\Gamma)$  the phase-space tangent vector field associated to an infinitesimal time evolution along the vector field  $t^a$  (which we allow to depend on the phase space point). Then  $\delta_t$  is Hamiltonian if there exists a functional  $H_t$  such that

$$\delta H_t = \Omega_M(\delta, \delta_t). \quad (58)$$

This requirement fixes a family of good energy formula since the Hamiltonian  $H_t$ , in presence of a boundary, acquires a surface term<sup>8</sup> and one can therefore define  $E_H^t$  as the surface term at  $H$  in the Hamiltonian—in addition to

---

<sup>8</sup>The volume (bulk) term in  $H_t$  is a linear combination of constraints, henceforth, they are absent in the covariant phase-space framework, since this consists only of solutions to the field equations.

the surface term at infinity representing the ADM energy. Remarkably, by means of the IH boundary conditions, the notion of isolated horizon energy  $E_H^t$  singled out by condition (58) automatically satisfies the first law of black hole mechanics, namely

$$\delta E_H^t = \frac{\kappa_{(t)}}{\kappa} \delta a_H + \Phi_{(t)} \delta Q_H + \text{other work terms}, \quad (59)$$

where we have put the explicit expression of the electromagnetic work term where  $\Phi_{(t)}$  is the electromagnetic potential (constant due to the IH boundary condition) and  $Q_H$  is the electric charge. The above equation implies  $\kappa_{(t)}$  and  $\Phi_{(t)}$  to be functions of the IH area  $a_H$  and charge  $Q_H$  alone.

In other words, the vector field  $\delta_t$  on  $\Gamma$  defined by the space-time evolution field  $t_a$  is Hamiltonian *if and only if* the first law (59) holds. The general treatment and derivation of the first law can be found in [48, 50].

Recently, a local first law for isolated horizons has been derived in [61], whose uniqueness can be proven once a local physical input is introduced. Interestingly, this allows to associate an energy to the horizon proportional to its area. This notion of area as energy could have important implications for statistical mechanical consideration of quantum IH (see, e.g., [62]).

## 5 Quantization

Isolated horizons—defined in Section 2—are the main ingredient we will use for the definition of a quantum black hole. Although very interesting developments [63, 64] have been recently carried out towards the characterization of black holes within the full quantum theory, we will follow here the well established, but somehow effective, approach based on starting from a classical space-time containing a black hole. Therefore, the goal is to quantize the sector of general relativity containing an isolated horizon as an internal boundary.

We will first present this quantization as it was originally carried out in [19] for the spherically symmetric case. However, we will only give an introductory description quantization procedure. For technical details and a thorough description the reader is referred to the original works [18, 19]. For computational simplicity, as described above, a gauge fixing was implemented that reduced the gauge group on the horizon from  $SU(2)$  to  $U(1)$ . That is, however, just a technical tool, not fundamental in the setup of this framework. Thus, in Section 5.2, we will present a more recent construction [41, 40, 42] in which this gauge fixing is avoided resulting in a quantum theory with a  $SU(2)$  gauge group on the horizon. These two approaches are expected to be fully equivalent; they should agree on all physical predictions. However, the different nature of the horizon constraints and number of free parameters in the theory within the two frameworks will have important

consequences in the entropy computation, as we are going to see in Section 6.

## 5.1 U(1) Quantization

The starting point is a space-time containing an inner boundary. As we have seen in the previous Section, it is not necessary to introduce a horizon boundary term in the action when working with vector-like variables. However, when passing to real Ashtekar-Barbero connection variables  $(A, \Sigma)$ , in order to have a conserved symplectic current, one needs to introduce a boundary term corresponding to the horizon in the symplectic structure. In the  $U(1)$  gauge fixed formulation, the resulting symplectic structure takes the form (56). As it can be seen from this expression, the boundary term corresponds to the symplectic structure of a  $U(1)$  Chern-Simons theory. This fact provides the main guidance on the strategy to quantize the system, namely to perform a separate quantization of the bulk and surface Hilbert spaces, where the surface Hilbert space will correspond to a quantum Chern-Simons theory on a (punctured) sphere.

From the structure of (56) it is possible to observe that we can split the phase-space in a bulk and a surface part. However, at the classical level, given a state in the bulk, the corresponding surface state is completely determined through continuity of the fields. The key point for the description of an entropy associated to the horizon is the fact that this no longer holds at the quantum level. Due to the distributional nature of quantum states, states in the horizon Hilbert space are not fully determined by the bulk. The horizon acquires independent degrees of freedom through the quantization process, and those are precisely the degrees of freedom giving rise to black hole entropy (we will come back to this point in Section 7).

### 5.1.1 Bulk Hilbert Space

Let us then start the quantization process by describing the bulk Hilbert space. The quantization of the bulk geometry follows the standard procedure of LQG [13] where one first considers (bulk) Hilbert spaces defined on a graph  $\gamma \subset M$  and then takes the projective limit containing the Hilbert spaces for arbitrary graphs.

As a result, geometry in the bulk is described by spin networks. However, due to the presence of the inner boundary, some of the spin network edges are not connected to vertices in the bulk, but they end at the horizon. These open ends of the spin network piercing the horizon  $H$ , denoted  $\gamma \cap H$ , are not connected to intertwiners and they acquire an additional label  $m$  (the spin projection) as a result of the non-gauge invariance at those points. Magnetic numbers  $m_i$  satisfy then the standard relation with the corresponding spin

labels  $j_i$  at each edge  $e_i$

$$m_i \in \{-j_i, -j_i + 1, \dots, j_i\}. \quad (60)$$

For a given set of points  $\mathcal{P}$  on the horizon  $H$  one can define the Hilbert space  $\mathcal{H}_M^{\mathcal{P}}$  as the space formed by all open spin networks with one edge finishing at each of the points of  $\mathcal{P}$ . We can assume, without loss of generality, that each puncture is connected to only one edge. Then, the total bulk Hilbert space can be written as the direct limit

$$\mathcal{H}_M = \lim_{\mathcal{P}} \mathcal{H}_M^{\mathcal{P}} \quad (61)$$

letting  $\mathcal{P}$  range over all finite sets of points in  $H$ . The quantum operator associated with  $\Sigma$  is

$$\hat{\Sigma}^i(x) = 8\pi\ell_p^2\beta \sum_{p \in \gamma \cap H} \delta(x, x_p) \hat{J}^i(p), \quad (62)$$

where  $[\hat{J}^i(p), \hat{J}^j(p)] = \epsilon^{ij}_k \hat{J}^k(p)$  at each  $p \in \gamma \cap H$ .

Another important operator that can be defined in  $\mathcal{H}_M^{\mathcal{P}}$  is the area of the horizon  $H$ . This operator is a particular case of the general area operator in LQG, in which edges only pierce the horizon from one side, and there are no edges lying on the surface. If we denote these spin network states by  $|\{j_p, m_p\}_1^N; \dots\rangle$ , where  $j_p$  and  $m_p$  are the spins and magnetic numbers labeling the  $N$  edges puncturing the horizon at points  $x_p$  (other labels are left implicit), the eigenvalues of the horizon area operator  $\hat{a}_H$  are

$$\hat{a}_H |\{j_p, m_p\}_1^N; \dots\rangle = 8\pi\beta\ell_p^2 \sum_{p=1}^N \sqrt{j_p(j_p + 1)} |\{j_p, m_p\}_1^N; \dots\rangle. \quad (63)$$

The bulk Hilbert space  $\mathcal{H}_M$  can be split into a direct sum of subspaces diagonalizing this operator. More precisely, for a given set of points  $\mathcal{P}$  and a set  $j = \{j_1, \dots, j_N\}$ , the space of all open spin networks with  $N$  edges ending at the points in  $\mathcal{P}$  and labeled by the set of spins in  $j$ , form the Hilbert space  $\mathcal{H}_M^{\mathcal{P}, j}$ . Then, the total volume Hilbert space  $\mathcal{H}_M$  can be written as

$$\mathcal{H}_M = \bigoplus_{\mathcal{P}, j} \mathcal{H}_M^{\mathcal{P}, j}. \quad (64)$$

This decomposition diagonalizes the horizon area operator  $\hat{a}_H$ . All the states in a subspace  $\mathcal{H}_M^{\mathcal{P}, j}$  correspond to the same area eigenvalue given by (63). Notice that throughout this construction the area of the horizon is taken as an operator acting on the volume Hilbert space. When referring to this area, we are thus referring to a geometrical property of the horizon  $H$  as a surface embedded in  $M$ .

An important issue that we will address later on is the gauge invariance of states in  $\mathcal{H}_M$ . As we saw above, there is a set of points  $\mathcal{P}$  whose gauge invariance is not established since they are associated to open edges of the spin network, not connected to an intertwiner. For the study of this gauge invariance it will be convenient to split the volume Hilbert space in a different way. Instead of using the above decomposition, that diagonalizes the geometric operators, it will be interesting to use a decomposition in terms of the spin projections  $m$ , that as we will see below diagonalizes the action of the gauge transformations. Taking this into account, one can consider the Hilbert subspace corresponding to spin networks piercing the horizon in a given set of points  $\mathcal{P}$  with spin projections  $m = \{m_1, \dots, m_N\}$ . Then, the volume Hilbert space can be written as

$$\mathcal{H}_M = \bigoplus_{\mathcal{P}, m} \mathcal{H}_M^{\mathcal{P}, m}. \quad (65)$$

As we will see, this decomposition turns out to be very convenient also for the computation of entropy.

### 5.1.2 Surface Hilbert Space

For quantizing the surface space, we start with a phase-space endowed with the symplectic structure of a Chern-Simons theory. Our phase-space is then made up of flat connections. However, through the quantization procedure, and due to the boundary conditions, those points where the spin network of the bulk pierce the horizon, behave as topological defects for the Chern-Simons theory. This gives rise to non-trivial degrees of freedom that will correspond to the horizon entropy. What is to be quantized, then, is a Chern-Simons theory over a punctured sphere. This quantization is carried out following a geometric quantization procedure, that we will not present in detail here (see [19] for details). After this, what is left is a Hilbert space formed by flat connections except at the punctures, where conical singularities of curvature occur. This distributional curvature concentrated at each puncture can be quantified as the angle deficit obtained when computing the holonomy of a path winding around the corresponding puncture. Therefore, these holonomies will be appropriate operators to encode the horizon degrees of freedom and then to characterize the quantum states. Furthermore, the angle deficits are quantized. The corresponding holonomies are given by

$$\hat{h}_i \Psi_{\mathcal{P}, a} = e^{\frac{2\pi i a_i}{k}} \Psi_{\mathcal{P}, a} \quad (66)$$

where  $a_i \in \mathbb{Z}_k$  are integer numbers modulo  $k$  that label the angle deficit at each puncture, and  $k$  is the level of the Chern-Simons theory. Thus, a convenient way to represent the states of the surface Hilbert space  $\mathcal{H}_H$  is by characterizing a set of punctures over the surface, and the corresponding

set  $a = \{a_1, \dots, a_N\}$  of labels associated to them. This is precisely what the state  $\Psi_{\mathcal{P},a} \in \mathcal{H}_H$  means in the above expression.

Although we will not get in details of the geometric quantization process here, there is however an important step that we need to comment on, given its relevance importance for the entropy counting.

The first step of the quantization is to construct the quantum phase-space, consisting of generalized connections that are flat everywhere except at the punctures. One has to show, then, that this phase-space is compact. This is done by showing that, for a given set  $\mathcal{P}$  of  $N$  punctures, the corresponding phase-space  $\mathcal{X}^{\mathcal{P}}$  is diffeomorphic to a  $2(N - 1)$ -torus. It can also be shown that, in spite of the existence of punctures with the corresponding singularities of curvature (curvature can be seen now as distributional), the Chern-Simons symplectic form is still well defined.

The important point that we want to discuss is related to this demonstration of the phase-space  $\mathcal{X}^{\mathcal{P}}$  being diffeomorphic to a  $2(N - 1)$ -torus. During this process, an additional structure consisting, among other things, of an ordering of the punctures is introduced. This ordering is then required for the quantization procedure to be well defined. But such a structure is not invariant under the action of diffeomorphisms on the horizon surface. In fact, diffeomorphisms act transitively on this additional structure. Thus, starting from a given ordering, one can obtain any other possible one, just through the action of a diffeomorphism. As a consequence, these different orderings cannot be considered as different physical states, since they are just related by a diffeomorphism. In fact, once one ordering is chosen, whatever this is, the physical states contained in the corresponding space are exactly the same as the ones that would be obtained with any other ordering. Then, one does not have to care about the ordering that is given to the punctures. But it is very important to keep in mind that, regardless of what the ordering is, the punctures have to be ordered in order for the quantization procedure to be well defined. This is a crucial point, as it affects the statistical character of the punctures. As ordered points, punctures have to be regarded as distinguishable objects. This will have a major relevance in the counting of states leading to black hole entropy.

With all this, the standard procedure for geometric quantization of  $\mathcal{X}^{\mathcal{P}}$  can be carried out, by first endowing this phase space with a Kähler structure with the symplectic form is its imaginary part and then defining a holomorphic line bundle  $L$ . This line bundle, however, can only be well defined if the level of the Chern-Simons theory takes an integer value  $k \in \mathbb{Z}$ . Then, the Hilbert space  $\mathcal{H}^{\mathcal{P}}$  is formed by holomorphic Sections of the line bundle  $L$ . This Hilbert space contains all the geometries of the horizon  $H$  that are flat everywhere except at the set  $\mathcal{P}$  of punctures. As commented above, a convenient basis for such a Hilbert space  $\mathcal{H}^{\mathcal{P}}$  is given by the states  $\Psi_{\mathcal{P},a}$ , characterized by the list  $a$  of the corresponding  $a_i$  labels associated to each puncture.

There is an additional consideration that can be made at this point. The spherical topology of the (spatial slices of the) horizon imposes a restriction reflecting the fact that holonomies of the  $N$  punctures are not all independent. A path winding around all the punctures is contractible on a sphere, so the composition of the individual holonomies for all the punctures must be identity. This fact translates into a constraint on the labels  $a_i$  of the holonomies, the so-called *projection constraint*:

$$\sum_{i=1}^N a_i = 0. \quad (67)$$

This constraint will also play an important role on the entropy counting.

With this, for a given set  $\mathcal{P}$  of points in the horizon surface and a certain labeling  $a$  for this points, satisfying (67), we can define a sub-space  $\mathcal{H}_H^{\mathcal{P},a}$  of the surface Hilbert space. The total surface Hilbert space  $\mathcal{H}_H$  can be then written as a direct sum of subspaces  $\mathcal{H}_H^{\mathcal{P},a}$

$$\mathcal{H}_H = \bigoplus_{\mathcal{P},a} \mathcal{H}_H^{\mathcal{P},a} \quad (68)$$

where the sum ranges over all finite sets of punctures  $\mathcal{P}$  labeled with nonzero integer numbers  $a_i \in \mathbb{Z}_k$  that sum up to zero. It is important to impose the condition of  $a_i$  being nonzero elements, in order for the direct sum decomposition to be well defined. In fact, if one considers a state with a puncture  $p_0$  labeled with  $a_0 = 0$ , this would correspond to having no curvature for this puncture. Physically, this state would be the same as a state with  $N - 1$  punctures, all of them with nonzero values of  $a_i$ . In order for the subspaces  $\mathcal{H}_H^{\mathcal{P},a}$  to be disjoint, avoiding a double counting of states, we have to require the nonzero condition on the  $a_i$  labels.

### 5.1.3 Quantum Boundary Conditions

Once the volume  $\mathcal{H}_M$  and surface  $\mathcal{H}_H$  Hilbert spaces have been separately constructed, we now have to impose the necessary conditions for matching states in these two spaces. It is at this point that one requires the boundary to be an isolated horizon, precisely by imposing the boundary conditions derived in the previous Section. The analysis of classical isolated horizons [18] shows that the pull back of the  $SU(2)$  Ashtekar-Barbero connection to a (spatial Section of the) isolated horizon can be fully characterized by the value  $a_H$  of the horizon area and a  $U(1)$  connection. Therefore, we can perform a gauge fixing on the constraint (24) by projecting this equation on a fixed internal vector  $r$  on the sphere, as illustrated in Section 3.3.1. This would allow us to work with a  $U(1)$  Chern-Simons theory on the quantum horizon. This gauge fixing is, however, not a necessary step, and we will see

in next Section how the quantization can be carried out keeping the  $SU(2)$  freedom.

When we project equation (24) on an internal vector  $r_i$ , and express it in terms of real Ashtekar-Barbero variables, we obtain the boundary condition (55), namely

$$F_{ab} = -\frac{2\pi}{a_H} \sum_{\leftarrow ab}^i r_i, \quad (69)$$

where, in order to match the notation in [19], we have introduced  $F_{ab} = dW$ , the curvature of the  $U(1)$  connection  $W$  obtained by projecting the pull-back to the horizon of the spin connection  $\Gamma_a^i$  on  $r_i$  (see (54)).

We will promote this equation to a quantum operator equation and impose it on the states of our Hilbert spaces. However, only the exponentiated version of the operator  $\hat{F}$  is well defined on the horizon. Thus, we will take the exponentiated version of the operators and establish the boundary condition as

$$(1 \otimes \exp(i\hat{F}))\Psi = (\exp(-i\frac{2\pi\beta}{a_H} \hat{\mathcal{J}} \cdot r) \otimes 1)\Psi. \quad (70)$$

This equation relates an operator acting on the surface with an operator acting on the volume. The structure of this equation implies that we can obtain a basis  $\Psi_M \otimes \Psi_H$  of solutions such that  $\Psi_M$  and  $\Psi_H$  are eigenstates of  $\hat{\mathcal{J}} \cdot r$  and  $\exp(i\hat{F})$  respectively. In order for the boundary condition to be satisfied, the spectra of these two operators should coincide. There is no *a priori* reason why this should be even possible. However, the spectrum of the operator  $\hat{\mathcal{J}} \cdot r$  satisfies

$$(\hat{\mathcal{J}} \cdot r)\Psi_M = 8\pi\ell_{\mathbb{P}}^2 \sum_{i=1}^N m_i \delta^2(x, p_i) \eta \Psi_M, \quad (71)$$

where  $p_i \in \mathcal{P}$  are the finitely many points in which the spin network punctures the horizon and  $m_i$  are the corresponding spin projection labels that the edges acquire at those points. Then at each point of  $\mathcal{P}$  the operator  $\exp(-i\frac{2\pi\beta}{a_H} \hat{\mathcal{J}} \cdot r)$  takes eigenvalues given by

$$\exp(-\frac{2\pi i\beta}{a_H} (8\pi\ell_{\mathbb{P}}^2 m_i)). \quad (72)$$

On the other hand, the operator  $\exp(i\hat{F})$  acts as the holonomy around each of the punctures  $\mathcal{P}$ , so, at each of this points the operator has eigenvalues given by

$$\exp(\frac{2\pi i a_i}{k}), \quad (73)$$

where  $a_i$  are the corresponding integers modulo  $k$  labeling the angle deficits. It is easy to see that these two spectra do indeed coincide if the relation

$$2m_i = -a_i \pmod{k} \quad (74)$$

is imposed between  $m$  and  $a$  labels.

Then, by using the splitting of the volume and surface Hilbert spaces in terms of  $m$  and  $a$  labels respectively, we can construct the total kinematical Hilbert space, including the boundary conditions, as

$$\mathcal{H}_{kin} = \bigoplus_{\mathcal{P}, m, a: 2m = -a \pmod k} \mathcal{H}_M^{\mathcal{P}, m} \otimes \mathcal{H}_H^{\mathcal{P}, a}. \quad (75)$$

The boundary conditions imposed by the isolated horizon definition are, at the end of the day, codified in a simple relation between labels of surface and volume states. States satisfying this relation are the ones giving rise to the kinematical Hilbert space.

It remains to impose the set of constraints necessary to construct the physical Hilbert space. Some additional comments can be made at this point. First, it can be shown that states satisfying this boundary conditions are automatically gauge invariant. The action of gauge transformations on the horizon turns out to be implemented by the same operators involved in the isolated horizon boundary condition, in such a way that states satisfying (74) are gauge invariant. Thus, the role played by equation (74) is precisely to ensure gauge invariance at the punctures. Second, in order for states to be diffeomorphism invariant, the position of punctures on the horizon cannot be a physical quantity. States that only differ on the localization of the punctures on the horizon, are related by a diffeomorphism and correspond to the same physical state. Thus, only the number  $n$  of punctures is needed to characterize physical states, and no reference to their position on the horizon will be made as a consequence of imposing the diffeomorphism constraint. Finally, since the Hamiltonian constraint has no effect on the horizon (the smearing function vanishes on the *horizon* [18]), it will suffice to make the (mild) assumption that for any horizon state there is at least one compatible bulk state satisfying the Hamiltonian constraint. Therefore, with all this we have a well defined physical Hilbert space with states satisfying Quantum Einstein's equations.

## 5.2 SU(2) Quantization

Let us now go back to the fully  $SU(2)$  invariant framework and relax any symmetry assumption on the horizon. In Section 3.3 we have seen that, for generic distorted horizons, one has to introduce two new  $SU(2)$  connections on the boundary and the horizon degrees of freedom are then described in terms of a pair of Chern-Simons theories satisfying the constraints (48).

Now, following Witten's prescription to quantize the two Chern-Simons theories with punctures [65], we introduce:

$$\frac{k_{\pm}}{4\pi} F^i(A_{\sigma_{\pm}}) = J_{\pm}^i(p), \quad (76)$$

where the levels  $k_{\pm}$  are given in (49). If we do so, we can now rewrite the constraints as [42]

$$D^i(p) = J_b^i(p) + J_\gamma^i(p) + J_\sigma^i(p) = 0 \quad (77)$$

plus the constraint

$$C^i(p) = J_\gamma^i(p) - J_\sigma^i(p) + \alpha J^i(p) = 0, \quad (78)$$

where

$$\alpha \equiv \frac{(\sigma_-^2 + \sigma_+^2) + \frac{\alpha_H}{\pi}(2\Psi_2 + c)}{(\sigma_-^2 - \sigma_+^2)} \quad (79)$$

will have a precise definition in the quantum theory in terms of bulk and boundary operator, as clarified in the following.

In a similar fashion as in Section 5.1.2, we can now quantize the boundary theory following Witten's prescription. In fact, the Hilbert space of the boundary model is that of two Chern-Simons theories associated with a pair of spins  $(j_p^+, j_p^-)$  at each puncture. More precisely,

$$\mathcal{H}_H^{CS}(j_1^+ \cdots j_N^+) \otimes \mathcal{H}_H^{CS}(j_1^- \cdots j_N^-) \subset \text{Inv}(j_1^+ \otimes \cdots \otimes j_N^+) \otimes \text{Inv}(j_1^- \otimes \cdots \otimes j_N^-). \quad (80)$$


The Hilbert space of an  $SU(2)$  Chern-Simons theory with given punctures on the sphere can be thought of as the intertwiner space of the quantum deformation of  $SU(2)$  denoted  $U_q(\mathfrak{su}(2))$ . The inclusion symbol in the previous expression means that the later space is isomorphic to a subspace of classical  $SU(2)$  intertwiner space. This is due to the fact that, in this isomorphism, the spins associated to the Chern-Simons punctures cannot take all values allowed by the representation theory of  $SU(2)$ , but are restricted by the cut-off  $k/2$  related to the deformation parameter by  $q = \exp(\frac{i\pi}{k+2})$  where  $k$  is the Chern-Simons level.

The operators associated to  $J_+^i(p)$  and  $J_-^i(p)$  describe the spins of the pair of Chern-Simons defects at the punctures. They are observables of the boundary system with which the spins  $j_p^+$  and  $j_p^-$  are associated. The theory is topological which means in our case that non-trivial d.o.f. are only present at punctures. The operator associated to  $J^i(p)$ , on the other hand, corresponds to the LQG flux operator (62) coming from the bulk.

Therefore, the distorted IH is modeled by a pair of  $SU(2)$  quantum intertwiners with each edge of one intertwiner coupled to an edge of the other and a puncture coming from the bulk. In a graphical representation, we have, at each puncture,

With all this we can now impose (77) which, at a single puncture, requires invariance under  $SU(2)$  local transformations

$$\delta J_A^j = [\alpha_i D^i, J_A^j] = \epsilon^{ij} \alpha_i J_A^k,$$



where  $J_A = J_+, J_-, J$  respectively. Equivalently, the constraint (77) requires the quantum state to be proportional to the singlet state with zero total  $SU(2)$  charge: zero total angular momentum. More precisely, the quantum constraint  $\hat{D}^i(p) = 0$  simply requires that

$$\text{Inv}(j_p \otimes j_p^+ \otimes j_p^-) \neq \emptyset \quad (81)$$

at each puncture  $p$ .

In order to analyze the imposition of the constraint (78), we first need to study the nature of the constraint system formed by  $\hat{D}^i(p), \hat{C}^i(p)$ . The constraint algebra is<sup>9</sup>:

$$\begin{aligned} [\hat{C}^i(p), \hat{C}^j(p')] &= \epsilon^{ij}_k (\hat{J}_+^k(p) + \hat{J}_-^k(p) + \alpha^2 \hat{J}^k(p)) \delta_{pp'} \\ [\hat{C}^i(p), \hat{D}^j(p')] &= \epsilon^{ij}_k \hat{C}^k(p) \delta_{pp'} \\ [\hat{D}^i(p), \hat{D}^j(p')] &= \epsilon^{ij}_k \hat{D}^k(p) \delta_{pp'}, \end{aligned} \quad (82)$$

from which we see that, in the generic distorted case, the constraint  $\hat{C}^i(p)$  is not first class. At this point, it is important to recall that in the spherically symmetric case, where the horizon degrees of freedom are described by a single  $SU(2)$  Chern-Simons theory, the boundary constraint (51) is first class since it closes an  $SU(2)$  Lie algebra, as shown in [41]<sup>10</sup>. As a consequence, for Type I IH one can correctly impose all the constraints strongly and end up with a single quantum  $SU(2)$  intertwiner modeling the horizon. This means that, in this new description, we need to be able to implement the constraint  $\hat{C}^i(p) = 0$  strongly for Type I IH in order to end up with the same number of d.o.f.

<sup>9</sup>Even though in the classical analysis, in order to prove that the symplectic structure is conserved, one has to identify the  $\Sigma$  in the bulk with that on the boundary, in the quantization process the degrees of freedom of the  $\Sigma$  bulk are decoupled from those on the boundary encoded in  $J_+, J_-$ . That's the reason why at the quantum level  $\hat{J}^i$  commutes with  $\hat{J}_+^i, \hat{J}_-^i$ .

<sup>10</sup>This is no longer the case in the  $U(1)$  gauge fixed version of the theory since the set of constraints (55), in the quantum theory, is no longer first class due to the non-commutativity of  $\Sigma^i$  in LQG. We will come back in the next Section on the consequences of this fact in the entropy computation.

Therefore, let us now concentrate for a moment on the spherically symmetric case and see if, within this more general framework, it is possible to recover the picture of [41]; it turns out that the answer is in the affirmative and the contact with Type I IH theory will allow us to reduce from two to one the number of free parameters  $(\sigma_+, \sigma_-)$  entering the system description.

As we saw in Section 3.3.1, for a spherically symmetric IH we can replace in the expression for  $\alpha$  (79)  $\Psi_2$  and  $c$  with their constant classical values:  $\Psi_2 = -\frac{2\pi}{a_H}$  and  $c = \frac{2\pi}{a_H}$ . In this way, the function  $\alpha$  becomes constant and we can use the freedom in the parameters  $\sigma_+, \sigma_-$  to set  $\alpha^2 = 1$ . If we do so, the algebra of constraints (82) now closes, reproducing a  $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$  local isometry. The previous analysis implies that we can impose spherical symmetry strongly if and only if  $(\sigma_-^2 + \sigma_+^2) \pm (\sigma_-^2 - \sigma_+^2) = 2$ . The solutions to the two branches of the previous equation are:  $\sigma_+^2 = 1$  and  $\sigma_-$  arbitrary (for the plus branch  $\alpha = 1$ ), and  $\sigma_-^2 = 1$  and  $\sigma_+$  arbitrary (for the minus branch  $\alpha = -1$ ). The two cases  $\alpha = \pm 1$  can be shown to be completely analogous and simply amount to switch the two indices ‘+’ and ‘-’. Henceforth, in the following we choose to fix  $\sigma_+^2 = 1$  and keep  $\sigma_-$  free. With this choice, the Chern-Simons levels  $k_{\pm}$  become

$$k_- = -k_+ = \frac{a_H}{4\pi\ell_p^2\beta(1 - \sigma_-^2)}. \quad (83)$$

If we now rewrite the constraints (77-78) as  $\hat{C}_{\pm}^i(p) \equiv (\hat{D}^i(p) \pm \hat{C}^i(p))/2$ , the algebra (82) becomes

$$\begin{aligned} [C_{\pm}^i(p), C_{\pm}^j(p')] &= \epsilon^{ij}_k C_{\pm}^k(p) \delta_{pp'} \\ [C_{\pm}^i(p), C_{\mp}^j(p')] &= 0 \end{aligned} \quad (84)$$

and we can impose  $C_{\pm}^i = 0$  strongly by setting the boundary spins  $j_p^- = 0$  and  $j_p^+ = j_p$ . In this way the Hilbert space of generic static isolated horizons  $\mathcal{H}_H^{CS}(j_1^+ \cdots j_n^+) \otimes \mathcal{H}_H^{CS}(j_1^- \cdots j_n^-)$  (restricted only by the condition (81)) reduces, for Type I IH, to

$$\mathcal{H}_H^{CS}(j_1 \cdots j_n) \subset \text{Inv}(j_1 \otimes \cdots \otimes j_n), \quad (85)$$

in complete agreement with the analysis of [41]. Moreover, the level (83) for the single  $SU(2)$  Chern-Simons theory left on the boundary exactly matches the value found in [41] for Type I IH.

This correspondence works also at the classical level. We have seen that spherical symmetry implies  $C_- = 0$  which, according to eq. (76), requires

$$F(A_{\sigma_-}) = 0; \quad (86)$$

as the horizon  $H$  is simply connected, this implies that  $A_{\sigma_-} = gdg^{-1}$ , *i.e.*, pure gauge. Therefore, the non-trivial degrees of freedom of the Type I isolated horizon are described by a single Chern-Simons theory with connection

$A_{\sigma_+}$  and constraint  $C_+ = 0$  equivalent to

$$\frac{k}{4\pi}F(A_{\sigma_+}) = \Sigma^i, \quad (87)$$

in complete classical correspondence with the treatment of [41].

Therefore, the requirement that spherical symmetry can be imposed strongly reduces the two-parameter family of models of a distorted horizon to a one-parameter one. Recall that this one-parameter ambiguity, also present in the spherical isolated horizon case, is a feature proper of the  $SU(2)$  treatment while is absent in the  $U(1)$  set-up.

We can now go back to generic distorted case and fix one of the two free parameters  $\sigma_+, \sigma_-$  in the general case to the value obtained from the requirement of the constraints to close a Lie algebra for the case of spherically symmetric IH. Namely, from now on, choosing the plus branch  $\alpha = 1$ , we will set  $\sigma_+^2 = 1$  and keep  $\sigma_-$  as the only free parameter. In the arbitrarily distorted case,  $\alpha$  is no more a constant and, therefore, the constraint algebra (82) doesn't close any more. This means that imposing the six constraints strongly is a far too strong requirement that risks to kill relevant physical degrees of freedom. Henceforth, in order to deal with the constraint  $\hat{C}^i(p) = 0$ , one has to introduce alternative techniques which allow to impose it weakly. A natural way of doing so consists of imposing  $\hat{C}^i(p) = 0$  strongly in the semi-classical limit, i.e. for large spins. In order to see what this implies, let us first notice that—using the closure constraint  $\hat{D}^i(p) = 0$ , which we do impose strongly as they are first class—the constraint  $\hat{C}^i(p)$  can be written in the following form

$$\hat{C}^i(p) \approx \hat{J}_+^i(p) - \hat{J}_-^i(p) - \alpha(\hat{J}_-^i(p) + \hat{J}_+^i(p)) = 0. \quad (88)$$

If we interpret for a moment the previous constraint classically, we see that it implies that the vectors  $J_+^i(p)$ ,  $J_-^i(p)$ , and (through  $D^i(p) = 0$ )  $J^i(p)$  are parallel. In order to derive this condition more rigorously, we are now going to introduce the master constraint technique [66]. More precisely, one can replace the constraint  $\hat{C}^i(p) = 0$  with the equivalent *master* constraint  $\hat{C}^2(p) = 0$ , which now commutes with  $\hat{D}^i(p) = 0$ . Trying to impose this master constraint strongly, one would find that the only states in the kernel of  $\hat{C}^i(p)$  are spherically symmetric states. Namely, the condition for  $\hat{C}^2(p)$  to vanish can be expressed as the restriction

$$\hat{J}_-^2(p)\hat{J}_+^2(p) - (\hat{J}_+(p) \cdot \hat{J}_-(p))^2 = 0, \quad (89)$$

which is equivalent to the vanishing of the quantum angle between  $\hat{J}_+^i(p)$  and  $\hat{J}_-^i(p)$ . The only strict solutions of that constraint are  $\hat{J}_-^i(p) = 0$  or  $\hat{J}_+^i(p) = 0$  which give  $\alpha = \pm 1$  and hence spherically symmetric states only. We can relax the previous constraint by requiring eq. (89) to hold only in the

large spin limit. Since the imposition of the closure constraint  $\hat{D}^i(p) = 0$  tells us that, at each puncture, we can decompose the boundary Hilbert space according to  $V_\rho = V_{j^+} \otimes V_{j^-} = \bigoplus_{j=|j^+-j^-|}^{j^++j^-} V_j$ , where  $V_j$  is the Irrep associated to the puncture coming from the bulk, the requirement of (89) to hold in the large spin limit amounts to select only the lowest and highest weight Irreps in the previous boundary Hilbert space decomposition. Namely, the weak imposition of  $\hat{C}^i(p) = 0$  correspond to the restriction

$$j = \begin{cases} j^+ + j^- \\ |j^+ - j^-|. \end{cases} \quad (90)$$

All this implies that it is consistent to take

$$\hat{\alpha} \equiv \frac{\hat{J}_+^2(p) - \hat{J}_-^2(p)}{\hat{J}^2(p)} \quad (91)$$

as a definition of the quantum operator associated to the horizon distortion degrees of freedom (79). The spins restriction (90) implies that the eigenvalues of the operator (91) are divided into the two sectors

$$|\alpha| \begin{cases} < 1 & \text{for } j = j^+ + j^- \\ > 1 & \text{for } j = |j^+ - j^-|, \end{cases} \quad (92)$$

with the eigenvalue  $|\alpha| = 1$  corresponding to spherically symmetric configurations. Notice that, for  $j = |j^+ - j^-|$ , the case where the two boundary punctures have the same spin, i.e.  $j_+ = j_-$ , is not allowed since it would give a  $j = 0$  for the bulk puncture and therefore excluded from the entropy counting.

Notice also that  $\hat{\alpha}$  defined in (91) commutes with all the observables in the boundary system; thus, the quantity  $2\Psi_2 + c$  remains ‘classical’ in this sense in agreement with the assumptions used for the construction of the phase-space of our system and the operator associated to it has no fluctuations in the Hilbert space of the distorted horizon.

Eq. (91) represents a well defined expression for an operator encoding the degrees of freedom of distortion; its eigenvalues are determined by the spins associated to the bulk and horizon punctures and they characterize the distorted configurations which will contribute to the entropy calculation. More precisely, the sum over the bulk and horizon spins performed (see next Section) in the state counting corresponds to the sum over the allowed distorted configurations of the model. In this sense, we can trace back the horizon entropy to the counting of the boundary geometry degrees of freedom.

## 6 Entropy Computation

Once we have described the quantization procedure and the resulting horizon and bulk Hilbert spaces, we want to compute the entropy associated to such

a black hole. As commented above, we consider the approach where only horizon degrees of freedom contribute to the black hole entropy. Hence, we need to trace out the degrees of freedom corresponding to the bulk. We construct the density matrix  $\rho_{bh}$  for the system and assume a maximally mixed state. In this way, the entropy of the horizon will be given by

$$S_{bh} = -Tr(\rho_{bh} \ln \rho_{bh}).$$

From standard statistical mechanics, we know that this is equivalent to  $S_{bh} = \ln N_{bh}$ , where  $N_{bh}$  is the total number of states in the horizon Hilbert space  $\mathcal{H}_H$ . Computing this number is the main goal of the rest of this Section, and in general, of the black hole entropy computation in LQG.

Thus, after all the formal quantization and setup of the framework, the main problem we are faced with, in order to obtain the behavior of the entropy of a black hole in loop quantum gravity, can be expressed as a purely combinatorial problem. In the following we state this combinatorial problem in a precise way.

At this point, it is important to comment on the different imposition of the constraints in the  $U(1)$  and  $SU(2)$  set-up. As already noted above, the set of constraints (55) in the  $U(1)$  formulation are no longer first class in the quantum theory due to the non-commutativity of  $\Sigma^i$  in LQG. Therefore, in the original derivation [18, 19] of the model, spherical symmetry is imposed already at the classical level. In this case, one considers a  $U(1)$  Chern-Simons theory with a level that scales with the macroscopic classical area  $k \propto a_H$ . This makes the state-counting (necessary for the computation of the entropy) a combinatorial problem which can be entirely formulated in terms of the representation theory of the classical group  $U(1)$ : for practical purposes one can take  $k = \infty$  from the starting point [25, 26, 35].

A striking result of these calculations is, besides the recovering of the leading term proportional to the horizon area, the appearance of logarithmic corrections in the Bekenstein-Hawking area law. In a first moment, these logarithmic corrections to the formula for black hole entropy in the loop quantum gravity literature were thought to be of the (universal) form  $\Delta S = -1/2 \log(a_H/\ell_p^2)$  [27]. However, in [67] it was pointed out that, assuming  $SU(2)$  gauge symmetry of the isolated horizon system, the counting should be modified leading to corrections of the form  $\Delta S = -3/2 \log(a_H/\ell_p^2)$ . This suggestion revealed to be particularly interesting since it would eliminate the apparent tension with other approaches to entropy calculation. In particular, the result of [67] is in complete agreement with the seemingly very general treatment (which includes the string theory calculations) proposed by Carlip [68], in which logarithmic corrections with a constant factor  $-3/2$  also appear<sup>11</sup>—see also [69] for an interesting relation between black

---

<sup>11</sup>See Section 9 for more details on the connections between Conformal Field Theory and the LQG description of the horizon theory.

hole thermodynamics and polymer physics in which a logarithmic correction with the same numerical coefficient is derived.

From this perspective, the necessity of an  $SU(2)$  gauge invariant formulation comes from the requirement that the isolated horizon quantum constraints be consistently imposed in the quantum theory, leading to the correct set of admissible states—in [40] it was suggested that the  $U(1)$  treatment leads to an artificially larger entropy due to the fact that some of the second class constraints arising from the  $SU(2)$ -to- $U(1)$  gauge fixing can only be imposed weakly<sup>12</sup>.

This observation, together with the basic conceptual ideas contained in the pioneering works [15]-[39], motivated the more recent derivation, performed in [40, 41, 42], of the horizon theory preserving the full  $SU(2)$  boundary symmetry. However, the  $SU(2)$  formulation is not unique as there is a one-parameter family of classically equivalent  $SU(2)$  connections parametrizations of the horizon degrees of freedom. More precisely, in the passage from Palatini-like variables to connection variables, that is necessary for the description of the horizon degrees of freedom in terms of Chern-Simons theory (central for the quantization), an ambiguity parameter arises, as shown in the previous section. This is completely analogous to the situation in the bulk where the Barbero-Immirzi parameter reflects an ambiguity in the choice of  $SU(2)$  variables in the passage from Palatini variables to Ashtekar-Barbero connections (central for the quantization in the loop quantum gravity approach). In the case of the parametrization of the isolated horizon degrees of freedom, this ambiguity can be encoded in the value of the Chern-Simons level  $k$ , which, in addition to the Barbero-Immirzi parameter, becomes an independent free parameter of the classical formulation of the isolated horizon-bulk system.

Therefore, it is no longer natural (nor necessary) to take  $k \propto a_H$ . On the contrary, it seems more natural to exploit the existence of this ambiguity by letting the Chern-Simons level be arbitrary. More precisely, we can reabsorb in the free parameter  $\sigma$  the dependence on  $a_H$  and thus take  $k \in \mathbb{N}$  as an arbitrary input in the construction of the effective theory describing the phase-space of IH. In this way, the  $SU(2)$  classical representation theory involved in previous calculations should be replaced by the representation theory of the quantum group  $U_q(su(2))$  with  $q$  a non-trivial root of unity [70]. Thus quantum group corrections become central for the state-counting problem.

The advantages of this paradigm shift introduced in [42] are that, on the one hand, it gives a theory which is independent of any macroscopic parameter—eliminating in this simple way the tension present in the old treatment associated to the natural question: *why should the fundamental quantum excitations responsible for black hole entropy know about the*

---

<sup>12</sup>Namely, in [19], for the last two constraints of (55), one has  $\langle \Sigma^i x_i \rangle = \langle \Sigma^i y_i \rangle = 0$ .

*macroscopic area of the black hole?*—on the other hand, compatibility with the area law will (as shown below) only fix the relationship between the level  $k$  and the Barbero-Immirzi parameter  $\beta$ ; thus no longer constraining the latter to a specific numerical value.

In the first part of this Section, we are going to present the powerful methods that have been developed for the resolution of the counting problem in the  $k = \infty$  case involving the  $U(1)$  classical representation theory [35, 36, 37, 32] (for a generalization to the  $SU(2)$  Lie group see [43]). In the second part, we present the finite  $k$  counting problem by means of simple asymptotic methods introduced in [44] and inspired by a combination of ideas stemming from different calculations in the literature [28, 67, 71]. This second part involves the quantum group  $U_q(su(2))$  representation theory and follows a less rigorous and more physical approach; perhaps, the more sophisticated techniques developed in the infinite  $k$  case are generalizable to the finite  $k$  case.

## 6.1 The Infinite $k$ Counting

As seen in Section 5.1, there are three sets of labels taking part on the description of the horizon-bulk quantum system. On the one hand, there are integer numbers  $a$  labeling the states on the surface Hilbert space. Corresponding to the bulk Hilbert space, and associated with each edge of the spin network piercing the horizon, there are two labels,  $j$  and  $m$ , that satisfy the standard angular momentum relations.  $j$  characterizes a  $SU(2)$  irreducible representation associated to the corresponding spin network edge, while  $m$  is the associated magnetic moment, therefore satisfying  $m \in \{-j, -j + 1, \dots, j\}$ . On the other hand, we have two constraints on them. The first constraint is the area of the horizon, and restricts the possible sets of spins  $j$ .

$$A(\vec{j}) = 8\pi\beta\ell_p^2 \sum_{i=1}^N \sqrt{j_i(j_i + 1)}. \quad (93)$$

The second is the projection constraint, that restricts the allowed configurations of  $a$  labels.

$$\sum_{i=1}^N a_i = 0. \quad (94)$$

Now, in principle, since we want to account only for the degrees of freedom intrinsic to the horizon, we should only be counting configurations labelled by  $a$ -numbers. However, the area constraint acts on labels  $j$ , and we still need to take it into account. Since we do not want to count degrees of freedom corresponding to the bulk, we need to find a way of translating the area constraint to the horizon states. Fortunately, we can make use of the relation (60) between  $j$  and  $m$  labels and also the isolated horizon

boundary condition (74) relating  $m$  and  $a$  labels. Noting this, a consistent way of posing the combinatorial problem was given in [25]:

$N_{bh}(A)$  is 1 plus the number of all the finite, arbitrarily long, sequences  $\vec{m}$  of non-zero half-integers, such that the equality

$$\sum_{i=1}^N m_i = 0. \quad (95)$$

and the inequality

$$8\pi\beta\ell_{\text{P}}^2 \sum_{i=1}^N \sqrt{|m_i|(|m_i| + 1)} \leq a_H. \quad (96)$$

are satisfied.

This problem was first solved in [26] for the large area limit approximation. An exact computational solution for the low area regime was later carried out in [29, 30], showing for the first time the effective discretization of black hole entropy in loop quantum gravity. What we are going to present in what follows is an exact analytical solution for this combinatorial problem, as it was performed later in [35]. This analytical exact solution has interest on its own, but it is also the point of departure for an asymptotic study of the behavior of entropy. It will allow to obtain closed analytical expressions for the behavior of entropy that can be used afterwards as the starting point for the asymptotic analysis. In order to solve this combinatorial problem we are going to use the following strategy (for a thorough exposition of this procedure see [37]):

- In first place, given a value  $A$  of area, we will compute all sets of integer positive numbers  $|m_i|$  such that the following *equality* is satisfied

$$\sum_{i=1}^N \sqrt{|m_i|(|m_i| + 1)} = \frac{a_H}{8\pi\beta\ell_{\text{P}}^2}. \quad (97)$$

This is equivalent to give a complete characterization of the horizon area spectrum in loop quantum gravity.

- For each set of  $|m_i|$  numbers we will introduce a factor accounting for all possible different ways of ordering them over the distinguishable punctures.
- Then, for each set of  $|m_i|$  numbers, we will compute all different ways of assigning signs to them in such a way that the projection constraint (95) is satisfied, thus getting the number of all possible  $\vec{m}$  sequences satisfying (95) and (97). We will call this quantity  $d_{DL}(A)$ .

- By adding up the degeneracy  $d_{DL}(A)$  obtained from the above steps for all values  $A$  of area lower than the horizon area  $a_H$  the complete solution to the combinatorial problem is obtained.

### 6.1.1 Area Spectrum Characterization

The first problem that we want to address is the characterization of the values belonging to the spectrum of the horizon area operator. In other words, the first question that we want to consider is: Given  $a_H \in \mathbb{R}$ , when does it belong to the spectrum of the area? Again, in order to simplify the algebra and work with integer numbers we will make use of the labels  $s_i$  defined as  $|m_i| = s_i/2$ , so that the area eigenvalues become

$$a_H = \sum_{i=1}^N \sqrt{(s_i + 1)^2 - 1} = \sum_{s=1}^{s_{\max}} n_s \sqrt{(s + 1)^2 - 1}. \quad (98)$$

Here we have chosen units such that  $4\pi\beta\ell_P^2 = 1$ , and the  $n_s$  (satisfying  $n_1 + \dots + n_{s_{\max}} = N$ ) denote the number of punctures corresponding to edges carrying spin  $s/2$ .

In order to answer this question, there is an important observation that we can make. Given any number  $\sqrt{(s + 1)^2 - 1}$ , one can always write it as the product of an integer  $q$  and the square root of a square-free positive integer number  $\sqrt{p}$  (SRSFN). A square-free number is an integer number whose prime factor decomposition contains no squares. Then, by using the prime factor decomposition of  $(s + 1)^2 - 1$  and factoring all the squares in it out the square root, one can always get the above structure. Hence, with our choice of units, every single area eigenvalue can be written as a linear combination, with integer coefficients, of SRSFN's. Only this integer linear combinations of SRSFN's  $\sum_I q_I \sqrt{p_I}$  can appear in the area spectrum. From now on, we will use these linear combinations to refer to the values of area. Then, the questions now are:

- First, given a linear combination  $\sum_I q_I \sqrt{p_I}$  of SRSFN's  $p_I$  with integer coefficients  $q_I$ , when does it correspond to an eigenvalue of the area operator?
- Then, if the answer is in the affirmative, what are the permissible choices of  $s$  and  $n_s$  compatible with this value for the area?

Answering to these questions is equivalent to giving a full characterization of the horizon area spectrum in LQG.

At this point, there is another important observation that we may do. The square roots of square free numbers are linearly independent over the rational numbers (and, hence, over the integers) i.e.,  $q_1 \sqrt{p_1} + \dots + q_r \sqrt{p_r} = 0$ , with  $q_I \in \mathbb{Q}$  and  $p_I$  different square-free integers, implies that  $q_I = 0$  for

every  $I = 1, \dots, r$ . This can be easily checked for concrete choices of the  $p_I$  and can be proved in general (see for instance [72]). We will take advantage of this fact in the following.

In order to answer the two questions posed above we will proceed in the following way. Given an *integer* linear combination of SRSFN's  $\sum_{I=1}^r q_I \sqrt{p_I}$ , where  $q_I \in \mathbb{N}$ , we need to determine the values of the  $s$  and  $n_s$ , if any, that solve the equation

$$\sum_{s=1}^{s_{\max}} n_s \sqrt{(s+1)^2 - 1} = \sum_{I=1}^r q_I \sqrt{p_I}. \quad (99)$$

Each  $\sqrt{(s+1)^2 - 1}$  can be written as an integer times a SRSFN so the left hand side of (99) will also be a linear combination of SRSFN with coefficients given by integer linear combinations of the unknowns  $n_s$ .

We can start by solving a preliminary step: for a given square-free positive integer  $p_I$ , let us find the values of  $s$  satisfying

$$\sqrt{(s+1)^2 - 1} = y \sqrt{p_I}, \quad (100)$$

for some positive integer  $y$ . At this point, it is very interesting to note that solving this equation is equivalent to solving a very well known equation in number theory, the Pell equation  $x^2 - p_I y^2 = 1$  where the unknowns are  $x := k + 1$  and  $y$ . Equation (100) admits an infinite number of solutions  $(s_m^I, y_m^I)$ , where  $m \in \mathbb{N}$  (see, for instance, [73]). These can be obtained from the fundamental one  $(s_1^I, y_1^I)$  corresponding to the minimum, non-trivial, value of both  $s_m^I$  and  $y_m^I$ . They are given by the formula

$$s_m^I + 1 + y_m^I \sqrt{p_I} = (s_1^I + 1 + y_1^I \sqrt{p_I})^m. \quad (101)$$

The fundamental solution can be obtained by using continued fractions [73]. Tables of the fundamental solution for the smallest  $p_I$  can be found in standard references on number theory. As we can see both  $s_m^I$  and  $y_m^I$  grow exponentially in  $m$ .

By solving the Pell equation for all the different  $p_I$  we can rewrite (99) as

$$\sum_{I=1}^r \sum_{m=1}^{\infty} n_{s_m^I} y_m^I \sqrt{p_I} = \sum_{I=1}^r q_I \sqrt{p_I}. \quad (102)$$

Using the linear independence of the  $\sqrt{p_I}$ , the previous equation can be split into  $r$  different equations of the type

$$\sum_{m=1}^{\infty} y_m^I n_{s_m^I} = q_I, \quad I = 1, \dots, r. \quad (103)$$

Several comments are in order now.

- First, these are diophantine linear equations in the unknowns  $n_{s_m^I}$  with the solutions restricted to take non-negative values. They can be solved by standard algorithms (for example the Fröbenius method or techniques based on the use of Smith canonical forms). These are implemented in commercial symbolic computing packages.
- Second, although we have extended the sum in (103) to infinity it is actually finite because the  $y_m^I$  grow with  $m$  without bound.
- Third, for different values of  $I$  the equations (103) are written in terms of disjoint sets of unknowns. This means that they can be solved independently of each other –a very convenient fact when performing actual computations. Indeed, if  $(s_{m_1}^{I_1}, y_{m_1}^{I_1})$  and  $(s_{m_2}^{I_2}, y_{m_2}^{I_2})$  are solutions to the Pell equations associated to different square-free integers  $p_{I_1}$  and  $p_{I_2}$ , then  $s_{m_1}^{I_1}$  and  $s_{m_2}^{I_2}$  must be different. This can be easily proved by *reductio ad absurdum*.

It may happen that some of the equations in (103) admit no solutions. In this case  $\sum_{I=1}^r q_I \sqrt{p_I}$  does not belong to the horizon area spectrum. On the other hand, if all these equations do admit solutions, then the value  $\sum_{I=1}^r q_I \sqrt{p_I}$  belongs to the spectrum of the area operator, the numbers  $s_m^I$  tell us the spins involved, and the  $n_{s_m^I}$  count the number of times that the edges labeled by the spin  $s_m^I/2$  pierce the horizon. Furthermore, if some of the equations in (103) admit more than one solution, then the set of solutions to (102) can be obtained as the cartesian product of the sets of solutions to each single equation in (103). Each of the sets of pairs  $\{(s_m^I, n_{s_m^I})\}$  obtained from this cartesian product will define a *spin configuration*  $\{n_s\}_{s=1}^\infty$  compatible with the corresponding value of area. We will call  $\mathcal{C}(a_H)$  the set of all configurations  $\{n_s\}_{s=1}^\infty$  compatible, in the sense of expression (97), with a given value of area  $a_H$  (note that, although the sets  $\{n_s\}$  can be formally considered to contain infinitely many elements  $n_s$ , the area condition (97) forces  $n_s = 0$  for all  $s$  larger than a certain value  $s_{max}(a_H)$ , so for all practical purposes the sets  $\{n_s\} \in \mathcal{C}(a_H)$  can be considered as finite). The number of different quantum states associated to each of these  $\{n_s\}$  configurations is given by two degeneracy factors, namely, the one coming from re-orderings of the  $s_i$ -labels over the distinguishable punctures (we will now call this  $r$ -degeneracy,  $R(\{n_s\})$ ) and the other originating from all the different choices of  $m_i$ -labels satisfying (95), (this will be now called  $m$ -degeneracy,  $P(\{n_s\})$ ). The  $r$ -degeneracy is given by the standard combinatorial factor

$$R(\{n_s\}) = \frac{(\sum_s n_s)!}{\prod_s n_s!} \quad (104)$$

We are going to compute the other factor in next Section.

### 6.1.2 Generating Functions

Let us consider then the  $m$ -degeneracy. The problem that we have to solve reduces to: Given a set of (possibly equal) spin labels  $s_i$ ,  $i = 1, \dots, N$ , what are the different choices for the allowed  $m_i$  such that (95) is satisfied?

This problem reduces just to find all different sign assignments to the  $m_i$  numbers in such a way that the total sum of them is zero. This problem is equivalent to solving the following combinatorial problem (closely related to the so called *partition problem*): Given a set  $\mathcal{O} = \{s_1, \dots, s_N\}$  of  $N$  (possibly equal) natural numbers, how many different partitions of  $\mathcal{O}$  into two disjoint sets  $\mathcal{O}_1$  and  $\mathcal{O}_2$  such that  $\sum_{s \in \mathcal{O}_1} s = \sum_{s \in \mathcal{O}_2} s$  do exist? The answer to this question can be found in the literature (see, for example, [74] and references therein) and is the following

$$P_{DL}(\mathcal{O}) = \frac{2^N}{M} \sum_{r=0}^{M-1} \prod_{i=1}^N \cos(2\pi r s_i / M), \quad (105)$$

where  $M = 1 + \sum_{i=1}^N s_i$ . This expression can be seen to be zero if there are no solutions to the projection constraint.

There is, however, a very powerful alternative approach to solving this problem: the use of generating functions. This approach has been extensively studied in [36], and provides very useful analytical expressions that can be used, as we will see, to study the asymptotic behavior of entropy. In particular, for this precise problem of computing the  $m$ -degeneracy, a generating function was obtained that gives rise to the following expression

$$P_{DL}(\{n_s\}) = \frac{1}{2\pi} \int_0^{2\pi} d\theta \prod_s (2 \cos(s\theta))^{n_s}. \quad (106)$$

By multiplying this factor by the reordering factor (104) for each  $\{n_s\}_{s=1}^{s_{max}}$  configuration, and summing the corresponding result for all different configurations of  $n_s$  contained in  $\mathcal{C}(A)$ , we obtain the corresponding degeneracy  $d_{DL}(A)$ . Finally, adding up  $d_{DL}(A)$  for all values of area  $A \leq a_H$ , the total horizon degeneracy corresponding to an horizon with area  $a_H$  is obtained. Once more, this problem can be solved by the use of generating functions. For details on how to obtain a generating function to solve the whole combinatorial problem, and how to use it in order to obtain closed analytic expressions for the solution, we refer the reader to [36, 37].

Finally, with very slight modifications, this whole procedure can be also applied to the SU(2) case. This was done in [43], and we will also comment the results of this computations in next Subsection.

### 6.1.3 Computational Implementation and Analysis of the Results

There are several ways of obtaining the results to the combinatorial problem presented above. One can simplify the problem from the beginning by

introducing some approximations and computing in certain limits. This was done in [26] for the large area limit, for instance. One can also implement the detailed procedure we just presented, involving number theory and generating functions, in a computer, and perform the computations by running an algorithm. This was also done in [35], and the results offer some new insights that were not initially detected in the large area limit. Finally, one can also try to extend the *exact* computation to the asymptotic limit in an analytic way. This is a much harder problem, but it has also been studied in [38], and we will present this analysis in next subsection. Let us summarize the main results for the entropy of a black hole in loop quantum gravity in the  $k = \infty$  case.

### Linear behavior

In the first place, the most remarkable result is that the entropy shows a linear behavior as a function of area. This was already obtained in the initial asymptotic computations and later corroborated by the computational results. This result constitutes the main test for the whole framework, as it shows compatibility with the expected Bekenstein-Hawking result. The first order expression for the entropy as a function of area is given by

$$S_{bh}(a_H) = \frac{\beta_H}{4\beta\ell_P^2} a_H, \quad (107)$$

where  $\beta_H$  is a constant obtained from the counting.

As one can see, the entropy grows linearly with the area, but there is also a freedom on the proportionality coefficient, as it includes the free Barbero-Immirzi parameter  $\beta$ . Therefore, the LQG  $U(1)$  counting reproduces the Bekenstein-Hawking law

$$S_{bh}(a_H) = \frac{a_H}{4\ell_P^2}$$

by choosing the appropriate value of this parameter  $\beta = \beta_H$ . Within this framework, the entropy computation can be regarded as a way of fixing the value of the Barbero-Immirzi parameter in the theory.

However, the value of the Barbero-Immirzi parameter obtained by this procedure for the two frameworks we are considering, namely the  $U(1)$  and  $SU(2)$  quantizations, turns out to be slightly different. This is another hint of inconsistency with the expectation that both approaches, when seen as one just a *gauge fixed* version of the other, should yield the same results. While this point of view is surely commonly shared at the classical level, in the quantum theory things become more subtle and the equivalence between the two approaches is far from obvious. For instance, we have already discussed above how the gauge fixing has important effects on the nature of the boundary constraints at the quantum level, affecting their correct Dirac

implementation and, therefore, the restriction to the proper set of admissible states. We will come back to this issue in Section 7.

### Logarithmic correction

If one looks at the next order, the first correction to the linear behavior is a logarithmic correction. This was also computed in [26, 25] and later confirmed by computational analyses. As already pointed out at the beginning of the Section, the coefficient  $C$  of this logarithmic correction depends on details of the counting, and in particular differs again between the  $U(1)$  and the  $SU(2)$  approaches being  $C = 1/2$  in the former and  $C = 3/2$  in the latter case. The entropy is then given by the formula

$$S_{bh}(a_H) = \frac{\beta_H}{\beta} \frac{a_H}{4\ell_P^2} - C \ln \frac{a_H}{\ell_P^2} + O(a_H^0) \quad (108)$$

It is interesting to point out, however, that the logarithmic correction is independent of the value of the Barbero-Immirzi parameter.

### Effective discretization

The third result was first observed in [29, 30], and consists of an effective discretization of the entropy for microscopic black holes, as a result of the particular *band structure* showed by the black hole degeneracy spectrum. This effect was not obtained in the first asymptotic calculations, and it only became apparent when an exact computational algorithm was implemented. A very extensive analysis of this phenomenon has been carried out during the last few years. Some of these works can be found in the bibliography [35, 36, 37, 29, 30, 33, 31], and the interested reader is encouraged to take a look at them. Here we will present a brief review of the main features.

When one plots the results for the black hole degeneracy spectrum  $d_{DL}(A)$  obtained by implementing the procedure presented in the previous Section in a computer, the results look like plot 3.

In order to obtain the entropy  $S_{bh}(a_H)$ , this quantity still has to be “integrated” for all values of area between 0 and  $a_H$ . When that sum is performed, the result is as shown in plot 4.

This effect is a very precise manifestation of the intricate structure of the black hole spectrum in loop quantum gravity, and has many potential implications. In particular, it can be seen to be in agreement with the Bekenstein conjecture on the discretization of black hole entropy, as it was pointed out in [30]. Furthermore, the structure of the degeneracy spectrum could leave some traces on the Hawking radiation spectrum, making it possible to detect a loop quantum gravity imprint on hypothetical microscopic black hole observations. This has recently been considered, and we will review the results in Section 10.




Figure 3: The degeneracy  $d_{DL}$  obtained from the number-theoretical procedure for each single area eigenvalue (in Planck units) is plotted.




Figure 4: The  $S_{bh}$  obtained from the number-theoretical procedure (in Planck units) is plotted as a function of the horizon area.

#### 6.1.4 Large Area Asymptotics

One of the fundamental questions in view of the discrete behavior of entropy for microscopic black holes described in the previous Section is whether this behavior is also present for large black holes or not. It is not possible to extend the computations to the large area limit by using computers, as it was done to obtain the results in the previous Section. If one wants to study the behavior of this discretization effect, one has to find some alternative approach. This was done in [38]. We are going to present a brief review here of the approach followed there.

The idea is to make use of some powerful theorems on analytic combinatorics, in particular about limits of distributions. We have already seen that it is possible to obtain a generating function to solve the computation of entropy. But in order to study this effect, it would be useful to have a generating function for one single *peak* of degeneracy (one of the individual bands that repeats periodically on the degeneracy spectrum). If we are able to study the limit distribution generated by such a generating function, we can understand the behavior of the band structure in this limit.

At this point, some ideas pointed out in [33] come at hand. There, it was shown that there is a way of characterizing each single peak by means of a given parameter  $K$ , constructed in terms of spins and number of punctures of the horizon configurations. Introducing this parameter into the generating function in an appropriate way allowed to obtain a generating function describing points *only* within one peak of the degeneracy spectrum. A theorem from [75] can be applied to this kind of generating function, and it is therefore possible to show that the generated distribution approaches a Gaussian when the value of area (or equivalently the value of the parameter  $K$ ) tends to infinity. Furthermore, it is possible to compute analytically the behavior of the mean and the variance corresponding to that Gaussian. All this was done in [38] and we refer the reader there for more details. The result is that the “Gaussian” peaks in the spectrum are wider as the area increases, reaching a point where they totally overlap each other, washing out the discrete behavior of entropy and giving rise, eventually, to a smooth linear growth. A model was constructed, using a superposition of the Gaussians obtained from the analytical computations. It showed that within the range of validity of the approximation, the discrete behavior completely washes out.

The conclusion is, therefore, that the effective discretization of entropy is a microscopic effect, only valid for black holes in their last stages of evaporation, and it disappears in the large area limit of astronomical black holes<sup>13</sup>.

---

<sup>13</sup>A note of caution is needed here, since the approach used in [38] cannot completely exclude the possibility of a revival of the effect at larger areas. This possibility seems, however, rather remote.

## 6.2 The Finite $k$ Counting

As illustrated at the beginning of this Section, an alternative strategy to deal with the counting problem and viable *only* in the  $SU(2)$  approach is to use the one-parameter freedom, introduced by the passage to the  $SU(2)$  connection variables on the horizon, to make the Chern-Simons level an arbitrary parameter (i.e. independent of the horizon area  $a_H$ ). In this way, the (now *finite*) level  $k$  enters the description of the boundary theory playing a role analog to the Barbero-Immirzi parameter in the bulk theory. We are now going to exploit further this ‘equal footing’ treatment and show how only the existence of a given relationship between the two free parameters of the boundary and bulk theories allows us to recover the Bekenstein-Hawking area law, eliminating in this way the need of fixing the Barbero-Immirzi parameter to the specific numerical value  $\beta_H$ .

Since the theory on the horizon is associated to Chern-Simons theory with punctures, dealing with a finite value of the level  $k$ , we cannot neglect anymore the quantum group representation theory underlying the structure of the Chern-Simons Hilbert space. Henceforth, first we first introduce an integral formulation of the dimension of the Chern-Simons theory Hilbert space  $\mathcal{H}^{CS}$ , when the space is a punctured two-sphere, which appears to be convenient to compute black hole entropy. Extending the relation between Chern-Simons and random walk, investigated in the classical case (namely, when  $k$  becomes infinite) in [71], to the quantum case (i.e. for a finite  $k$ ), it can be shown that, for a set of  $p$  punctures, denoted by  $\ell \in [1, p]$ , each labeled by an unitary irreducible representation  $j_\ell$  of the quantum group  $U_q(su(2))$ , the dimension of the punctured 2-sphere Hilbert space can be expressed as [44]

$$N_k(\mathbf{d}) = \frac{1}{\pi} \int_0^{2\pi} d\theta \sin^2\left(\frac{\theta}{2}\right) \frac{\sin\left(\left(r + \frac{1}{2}\right)k\theta\right)}{\sin\frac{k\theta}{2}} \prod_{\ell=1}^p \frac{\sin\left(d_\ell \frac{\theta}{2}\right)}{\sin\frac{\theta}{2}}, \quad (109)$$

where the dimension  $d_\ell = 2j_\ell + 1$  of the  $j_\ell$ -representation is the same as in the classical theory (even though now we have the cut-off  $j_\ell \leq k/2$ ) and  $r \equiv \lfloor \sum_{\ell=1}^p (d_\ell - 1)/(2k) \rfloor$  (here  $\lfloor x \rfloor$  is the floor function). Notice that  $N_k(\mathbf{d})$  coincides with the classical formula when  $r = 0$ , i.e. when  $\sum_{\ell=1}^p (d_\ell - 1) < 2k$ .

Let us now briefly recall that the entropy of an IH is computed by the formula  $S = \text{tr}(\rho_{IH} \log \rho_{IH})$ , where the density matrix  $\rho_{IH}$  is obtained by tracing over the bulk d.o.f., while restricting to horizon states that are compatible with the macroscopic area parameter  $a$ . Assuming that there exists at least one solution of the bulk constraints for every admissible state on the boundary, the entropy is given by  $S = \log(N(a))$  where  $N(a)$  is the number of admissible horizon states. Henceforth, the entropy calculation problem boils down to the counting, in the large horizon area limit, of the dimension of the horizon Hilbert space. For a generic distorted IH, we have

seen in Section 5.2 that the Hilbert space takes the form (80), for which the dimension of each  $\mathcal{H}_H^{CS}$  is expressed by the formula (109).

Now, following the techniques introduced in [44], the entropy  $S(a) = \log N(a)$  of a distorted isolated horizon of macroscopic area

$$a = \frac{1}{2} \sum_{\ell=1}^p \sqrt{(d_\ell - 1)(d_\ell + 1)}, \quad (110)$$

where  $a \equiv a_H/8\pi\beta\ell_p^2$ , is defined from the number of states

$$\begin{aligned} N(a) &= \sum_{p=0}^{\infty} \sum_{\mathbf{d}} \delta\left(a - \frac{\sum_{\ell=1}^p \sqrt{(d_\ell - 1)(d_\ell + 1)}}{2}\right) \\ &\cdot \sum_{\mathbf{d}^+, \mathbf{d}^-}^{k+1} \left( \prod_{\ell=1}^p Y(j_\ell, j_\ell^+, j_\ell^-) \right) N_k(\mathbf{d}^+) N_k(\mathbf{d}^-), \end{aligned} \quad (111)$$

where the sums run over the families  $\mathbf{d}^\pm = (d_1^\pm, \dots, d_p^\pm)$ ,  $\mathbf{d} = (d_1, \dots, d_p)$  of representations dimensions associated with the boundary and bulk punctures  $j_\ell^\pm, j_\ell$ . In the previous expression, in order to implement the admissibility condition,  $Y_\ell \equiv Y(j_\ell, j_\ell^+, j_\ell^-) = 1$  if  $(j_\ell, j_\ell^+, j_\ell^-)$  satisfy the restriction (90) at each puncture, it vanishes otherwise. If one assumes that the number of states grows exponentially with the area, the study of the entropy for large  $a$  but finite  $k$  can be performed by means of the Laplace transform of  $N(a)$ , namely

$$\tilde{N}(s, t) = \int_0^\infty da e^{-as} a^{-t} N(a), \quad (112)$$

and the number theory and the complex analysis aspects illustrated in the previous Subsection are not necessary to get the main ideas and results. In fact, the Laplace transform technique allows us to study the leading and sub-leading terms in the asymptotic expansion of  $N(a)$ . More precisely, assuming an asymptotic behavior of the form

$$N(a) \sim e^{s_c a} a^{-(t_c+1)}, \quad (113)$$

at large  $a$ , one can obtain the critical exponents  $s_c$  and  $t_c$  by studying the convergence properties of the integral  $\tilde{N}(s, t)$ . This study has been done in [44]; here we just report the results.

In the spherically symmetric case, the critical exponent  $s_c$  is the unique solution of the equation

$$1 - \sum_{d=1}^k (d+1) e^{-\frac{s_c}{2} \sqrt{d(d+2)}} = 0, \quad (114)$$

which encodes the dependence of the Barbero-Immirzi parameter  $\beta$  on the level  $k$  as we required the recovering of the Bekenstein-Hawking area law for the leading term. In fact, one has

$$S = \log(N(a)) = \frac{a_H}{4\ell_p^2} + o(\log a_H) \quad (115)$$

as soon as  $s_c = 2\pi\beta$ . For increasing values of the level  $k$ , the solutions of equation (114) for  $\beta$  reach fast an asymptotic value which coincides, as expected, with the value  $\beta_H$  found in [43] when  $k \rightarrow \infty$ .

For the sub-leading term one finds the critical exponent  $t_c = 1/2$  from which

$$N(a) \sim e^{s_c a} a^{-3/2} \quad \text{for large } a. \quad (116)$$

Therefore, even if the finiteness of the level  $k$  affects the behavior of the leading term, it does not modify the sub-leading corrections when  $a$  is large. In that sense, the logarithmic corrections seem to be independent of the Barbero-Immirzi parameter even in the  $SU(2)$  spherically symmetric black hole, confirming the results of [67, 43] and they show the universal nature conjectured in [68].

In the distorted case, it can be shown that, for  $k$  large enough,  $s_c$  grows logarithmically with  $k$ , while the critical exponent of the logarithmic corrections is  $t_c = 2$ . Eq. (113) then shows that the model presented in Section 5.2 for static IH provides a leading order entropy recovering exactly Hawking's area law (plus logarithmic corrections)

$$S = \frac{a_H}{4\ell_p^2} - 3\log(a_H) \quad (117)$$

once the following relationship between the Barbero-Immirzi parameter  $\beta$  and the Chern-Simons level  $k$  holds

$$1 - \sum_d \sum_{d^\pm=0}^k Y(d^+ + 1)(d^- + 1)e^{-\pi\beta\sqrt{d(d+2)}} = 0. \quad (118)$$

The previous equation is obtained from the condition determining the value of the critical exponent  $s_c$ , as for (114) in the spherically symmetric case; its numerical solution, for the first integer values of  $k$ , is plotted in Figure 5. The plot shows that the Barbero-Immirzi parameter grows, for values of the level large enough, as

$$\beta_k = \sqrt{3}/\pi \log(k + 1) + \mathcal{O}(1), \quad (119)$$

where the constant  $\sqrt{3}/\pi$  is obtained from eq. (118) in the large  $k$  limit and assuming that all the bulk spins  $j$  are fixed to  $1/2$ .

The factor  $-3$  in front of the logarithmic corrections in the distorted case can be traced back to the fact that now, instead of a single  $SU(2)$  Chern-Simons theory describing the horizon d.o.f., we have two of them, as a consequence of any symmetry assumption relaxation.




Figure 5: In the figure we plotted the values of the Barbero-Immirzi parameter  $\beta_k$  as function of  $k \in \mathbb{N}$  for the first integers; the plot shows a logarithmic growth of the Barbero-Immirzi parameter with the level.

## 7 On the Nature of the Entropy Degrees of Freedom

After all this presentation of the black hole entropy calculation in LQG, both in its original  $U(1)$  derivation and in its more recent fully  $SU(2)$  invariant set-up, some questions arise naturally: What is, at the end, the nature of the degrees of freedom accounting for the black hole entropy? Or, in other words, what are these models really counting? Is there any difference in the identification of these degrees of freedom between the  $U(1)$  and the  $SU(2)$  frameworks?

Addressing these questions is the main goal of all the construction presented so far but the same answer is not always shared by all the community. We now want to try to clarify this point and present some different perspectives.

Let us start with the spherically symmetric case in its original  $U(1)$  formulation [18, 19]. In this case, at the classical level, the system is characterized by a single degree of freedom corresponding to the horizon macroscopic area. In fact, the classical boundary theory contains no independent states. Independent boundary states arise only at the quantum level since the quantum configuration space is larger than the classical one, as a consequence of the fact that the former admits distributional connections. More precisely, the classical configuration space  $\mathcal{A}$  of general relativity can be taken to consist of smooth  $SU(2)$  connections on the spatial 3-manifold  $M$ . Its completion  $\bar{\mathcal{A}}$  consists of ‘generalized’  $SU(2)$  connections and this is what represents the quantum configuration space since, in quantum field theories with local degrees of freedom, quantum states are functions of generalized fields which need not be continuous. In fact, holonomies of generalized connections are not required to vary smoothly with the path and, therefore,  $\bar{\mathcal{A}}$

turns out to be very large<sup>14</sup>.

In other words, in the classical theory, even though the symplectic structure contains also a Chern-Simons term for a connection on the internal boundary, the boundary connection does not represent new degrees of freedom, since it is determined by the limiting value of the connection on the bulk. At the quantum level though, states are functions of generalized connections fields which need not to be continuous. Therefore, the behavior of generalized connections on the boundary  $H$  can be quite independent of their behavior in the bulk; it follows that, at the quantum level, surface states are no longer determined by bulk states.

The boundary condition (24) is such that, given the value of the area  $a_H$ , the connection is unique up to gauge and diffeomorphisms. Henceforth, at the classical level, there are no true ‘configuration space’ degrees of freedom on the horizon. However, at the quantum level, when one first quantizes and then imposes the constraints, the horizon boundary condition becomes an operator restriction on the allowed quantum states. More precisely, both the boundary connection  $A$  and the flux field  $\Sigma$  are allowed to fluctuate but they do so respecting the quantum version of (24). Imposing this restriction leads to the appearance of Chern-Simons theory with punctures which has a finite number of states. This is the theory describing the geometry of the quantum horizon and accounting for its entropy.

From this point of view, one has one physical (classical) macrostate which corresponds to a large number of (quantum) microstates arising through quantization: It is the quantum theory that ‘multiplies’ the number of degrees of freedom.

The distorted case in the  $U(1)$  framework has been firstly treated in [54]. In that approach, the distortion degrees of freedom contained in the real part of the Weyl tensor component  $\Psi_2$  are encoded in the values of some geometric multipoles which provide a diffeomorphism invariant characterization of the horizon geometry. Thanks to the additional assumption of axisymmetry, the system is then mapped to a model equivalent to the Type I case if the horizon area and the multipole moments describing the amount of distortion are fixed classically. Therefore, for fixed area and multipoles, the boundary theory is still described in terms of a *fiducial* Type I  $U(1)$  connection, satisfying the boundary condition (54). In this way, the problem of quantization reduces to that of spherically symmetric IH and the mathematical construction of the physical Hilbert space presented in Section 5.1 can be taken over.

Taking the limit  $k \rightarrow \infty$ , one can associate an operator to the Weyl tensor component  $\Psi_2$  and the multipoles, whose eigenvalues can be expressed in terms of the classically fixed values of the area and the multipoles and the eigenvalues of the total area operator associated with  $H$ . In other words,

---

<sup>14</sup>Recall that the quantum configuration space  $\bar{\mathcal{A}}$  is constructed through projective limit of configuration spaces  $\mathcal{A}_g$  of  $SU(2)$  lattice gauge theory associated with a finite graph  $g$ .

even if classically fixed, the multipoles can have quantum fluctuations and these are dictated by the fluctuations in  $\hat{a}_H$  [54].

However, all this construction of quantum operators encoding Type II horizon quantum geometry is argued to be inessential to the entropy counting. This is due to the mapping to the equivalent Type I model and the observation that the counting of the number of states in the micro-canonical ensemble for which the horizon area and multipoles lie in a small interval around their classically fixed values is, in this approach, the same as in the spherically symmetric case. Hence, the horizon entropy is again given by (108) with the same value of  $\beta_H$  found in the Type I analysis and  $C = 1/2$ .

To summarize, the approach of [54] to incorporate distortion degrees of freedom consists of introducing an infinite set of multipoles to capture distortion and then define a Hamiltonian framework for the sector of general relativity consisting of space-times which admit an IH with fixed multipoles. The resulting phase-space is then mapped to one equivalent to a Type I IH in order to use the counting techniques developed for this simpler case.

Classically, the complete collection of multipole moments characterize an axisymmetric horizon geometry up to diffeomorphism. Fixing the values of the area and the multipoles classically allows to select a phase-space sector of the full classical one, corresponding to a given distorted intrinsic geometry and all the others related to that by a diffeomorphism. However, if one wants to take into account all possible kind of axisymmetric classical distortions, one would end up with a pile of different phase-space sectors, which cannot be related by a diffeomorphism. Each of these sectors would now have to be mapped to a different Type I model, naively leading, in this way, to an infinite entropy. The situation seems even worse if one takes into account also all the non-axisymmetric configurations.

This issue has recently been addressed in [76]. In this work, the authors relax the axisymmetry assumption in order to deal with generic horizon geometries. Remaining within the  $U(1)$  framework, they show how it is possible to quantize the *full* phase-space of *all* distorted IH of a given area without having to fix classically a sector corresponding to a particular horizon shape, with the resulting Hilbert space identical to that found previously in [19]. More precisely, they manage to extend the map to a spherically symmetric  $U(1)$  connection introduced in [54] to the generic distorted case, which, however, now becomes non-local. Then they argue that the boundary term in the symplectic structure for the full classical phase-space of all isolated horizons with given area can be expressed in terms this Type I connection and that all elements of the classical framework necessary for quantization in [19] are also present in this more general context. This leads to the *reinterpretation* of the quantization described in [19] as that of the full phase-space of generic isolated horizons. Even further, [76] claims that the physical Hilbert space as constructed in [19] does not incorporate spherical symmetry.

The point of view of [76] is, therefore, quite the opposite of the original

one expressed in [18, 19]. At the classical starting level one does not have any more a single degree of freedom (the value of the horizon area) which then gets multiplied by the quantization procedure. On the contrary, one starts with an infinite number of degrees of freedom, associated to all possible distortions of the intrinsic geometry, which would naively lead to the expectation of an infinite entropy. But now the quantum structure plays the inverse role by providing a cut-off to this infinite pile of degrees of freedom, rendering the entropy finite.

Reinterpreting the quantization in this way, it is natural to consider different horizon states in the Hilbert space to represent different quantum shapes of the horizon. This reinterpretation of the physical Hilbert space built in [19] reconciles the entropy calculation with the conceptual viewpoint that entropy arises by counting different microscopic shapes of the horizon intrinsic geometry, proposed in [39] and recently also investigated in [69].

The point of view of [76] is similar in spirit to the one adopted in [42], and described in Section 5.2, for the definition of a statistical mechanical ensemble accounting for the degrees of freedom of generic distorted  $SU(2)$  IH. In [42] no symmetry assumption is necessary either (Type I, Type II, and Type III horizons are all treated on equal footing), only staticity is a necessary condition for the dynamical system to be well defined. However, the approach of [42] differs from the previous works [54, 76] dealing with distorted IH in two main respects: first the treatment is  $SU(2)$  gauge invariant, avoiding in this way the difficulties found upon quantization in the gauge fixed  $U(1)$  formulation, and second, distortion is not *hidden* by the choice of a mapping to a canonical Type I connection. In particular, the degrees of freedom related to distortion are encoded in observables of the system which can be quantized and are explicitly counted in the entropy calculation. In this new treatment, as shown in Section 5.2, one can find the old Type I theory in the sense that when defining the statistical mechanical ensemble by fixing the macroscopic area  $a_H$  and imposing spherical symmetry, one gets an entropy consistent with the one in [41].

More precisely, as in [76], the starting point of [42] is again the full classical phase-space of all distorted IH and, avoiding the passage to a non-local Type I connection, both intrinsic and extrinsic<sup>15</sup> geometry degrees of freedom can be quantized, leading to the definition of the distortion operator (91). This operator has a discrete spectrum and its eigenvalues are bounded by the cut-off introduced by the *finite*<sup>16</sup> Chern-Simons level. Henceforth,

---

<sup>15</sup>Recall that, due to the more generic treatment required by inclusion of distortion, eq. (26), relating intrinsic and extrinsic curvatures, plays a central role in the construction of the conserved symplectic structure of the system and the curvature scalar  $c$  enters the definition (79) of  $\alpha$ .

<sup>16</sup>The passage to a finite level  $k$ , independent of the horizon area  $a_H$  [42], is a crucial step in the entropy calculation. In fact, if one keeps the linear growth of  $k$  with  $a_H$ , it can be shown that taking into account all the distorted d.o.f. leads to a violation of the area

even though at the classical level we had an infinite number of distortion degrees of freedom—encoded in all the possible (continuous) values of the real part of  $\Psi_2$  and of the curvature invariant  $c$ —the physical Hilbert space defined by (80) with the restrictions (81)-(90) provides a finite answer for the entropy (117) due to the cut-off introduced by the quantum group structure and the consistency with the area constraint (required by the gauge invariance condition (81)). This entropy can be directly traced back to the distorted horizon quantum theory, accounting for both intrinsic and extrinsic geometry degrees of freedom.

The analysis carried out in [76] and [42] surely provides a conceptually common (to both the  $U(1)$  and the  $SU(2)$  approaches) framework to understand the black hole entropy counting in LQG and try to answer coherently the questions raised at the beginning of the Section. Nevertheless, a deeper understanding of the relation between the two constructions seems necessary in order to have a clearer description of the distorted quantum geometry of the horizon. In this direction, it seems important to investigate further the role played by transverse fluxes operators (i.e. fluxes  $\hat{\Sigma}[T, f]$  through surfaces  $T$  intersecting  $H$  transversely) in the characterization of the horizon intrinsic geometry advocated in [76] and the properties of the distortion operator (91)<sup>17</sup>.

## 8 On the Role of the Barbero-Immirzi Parameter

As show in Section 6, one of the main success of the black hole entropy calculation in LQG is the recovering of the Bekenstein-Hawking area law for the leading term in the asymptotic large area limit. This result was first derived within the symmetry reduced  $U(1)$  model [18, 19], but left many people not fully satisfied, since it implied an ‘unpleasant’ constraint for the full quantum theory: In order to recover the numerical factor  $1/4$  in the Bekenstein-Hawking formula, one had to fix the value of the Barbero-Immirzi parameter, entering the calculation through the form of the area spectrum in LQG, to a specific numerical value, as shown in Section 6.1.3.

Let us recall that the Barbero-Immirzi parameter has been introduced in canonical non-perturbative quantum gravity to deal with the difficulties raised by the non-compactness of the gauge group in relation to the passage to Ashtekar self-dual variables. With the use of real connection variables the gauge group becomes compact (the connection is now  $SU(2)$  Lie algebra valued) and the formalism relevant at the diffeomorphism-invariant and background-independent level easier to handle. However, even if classi-

---

law—namely, one obtains a leading term for the entropy of the form  $S \approx a_H \log(a_H)$ .

<sup>17</sup>Recall that the definition of the distortion operator (91) is closely related to the vanishing, in the large spin limit, of the quantum angle between  $\hat{J}_+$  and  $\hat{J}_-$  [42].

cally irrelevant<sup>18</sup>, the Barbero-Immirzi parameter represents a quantization ambiguity in the kinematics of LQG; in fact, to different values of  $\beta$  corresponds, due to its appearance in the spectrum of geometric operators, different sectors of the quantum theory which cannot be related by unitary transformations [77]. Nevertheless, this  $\beta$  dependence would better drop out from any physical prediction of the theory.

In this sense, using Hawking semi-classical analysis on black hole radiation and entropy associated to it to fix the value of a parameter which plays a role only in the quantum theory doesn't seem very natural<sup>19</sup>. Moreover, this would put a very stringent constraint on the consistency of the theory and the determination of the same value from at least another application of LQG would be required to strengthen the validity of this point of view. In other words, the need to constrain  $\beta$  to a specific numerical value has so far been seen as the Achille's heel of the LQG computation, mostly from the perspective of alternative approaches to black hole entropy derivation but partly also within the community.

This issue has lately received a lot of attention again and it has, somehow, also motivated the definition of the fully  $SU(2)$  invariant formulation of IH quantization recently provided in [40, 41, 42]. As shown in Section 5.2, from this analysis it emerged that, when avoiding the symmetry reduction, an extra ambiguity parameter appears in the quantum theory describing the horizon degrees of freedom in terms of  $SU(2)$  connection variables. This new parameter in the boundary theory has exactly the same origin as the Barbero-Immirzi parameter in the bulk and, when exploiting this analogy in the entropy calculation, consistency with the Bekenstein-Hawking area law doesn't fix the value of  $\beta$  anymore. More precisely, as elucidated in Section 6.2, the new ambiguity can be encoded in the level of the Chern-Simons theory, describing the boundary degrees of freedom, to eliminate its dependence on the horizon area and render it a free input. In this way, the level is now finite and, by means of quantum groups representation theory, the leading order  $a_H/(4\ell_p^2)$  for generic distorted IH entropy can be recovered as long as  $\beta$  and  $k$  satisfy (118) [41]. This implies that consistency with the Hawking analysis now requires, in the semiclassical regime (i.e for large values of  $k$ ) the Chern-Simons level to grow logarithmically with the Barbero-Immirzi parameter according to (119).

Therefore, the numerical restriction on the value of  $\beta$  turns out to be just an artifact of the  $U(1)$  symmetry reduction. While it is encouraging

---

<sup>18</sup>In the classical theory the Barbero-Immirzi parameter introduces a symmetry which can be realized as a canonical transformation.

<sup>19</sup>See, for example, [78] where the authors apply the Wald [79] approach to compute entropy from Noether charges to the case of first order gravity with a negative cosmological constant and adding the Holst term. They show that the AdS-Schwarzschild black hole entropy obtained in this way from the semi-classical theory presents no Barbero-Immirzi parameter dependence.

that the  $SU(2)$  treatment contains a free parameter arising at the boundary from exactly the analogous reason as the Barbero-Immirzi parameter appears in the bulk parametrization of the phase space. This keeps open the possibility that dynamical considerations could lead to cancelation of both ambiguities producing Barbero-Immirzi parameter independent predictions. On a speculative level, this could be obtained, for example, by deriving the relation (119) through semi-classical considerations involving Schwarzschild near-horizon geometry.

Another possibility to eliminate the Barbero-Immirzi parameter dependence from the leading term in the entropy has also, very recently, appeared in [45]. In this work, the authors propose an alternative analysis of black hole entropy in the LQG approach which gives a result in agreement with Hawking's semiclassical analysis for all values of the Barbero-Immirzi parameter. The key ingredient consists of a modification of the first law of black hole mechanics by taking into account the underlying quantum geometry description of the black hole horizon. More precisely, [45] introduces a *quantum hair* for the black hole, related to the number  $N$  of topological defects in the quantum isolated horizon and proportional to some chemical potential  $\mu$ , and exploits the consequences of having a minimal length in the quantum theory of the order of the Planck scale. By doing so, they show how, from the statistical mechanics of the basic quantum excitations of IH in LQG, the leading order  $a_H/(4\ell_p^2)$  for the entropy can be recovered, with all the dependence on the Barbero-Immirzi parameter shifted on a chemical potential correction term which vanishes as soon as the relation (118) holds. In this way, the Barbero-Immirzi parameter appears in an additive correction to the semiclassical expression. This additive term is the quantum correction to the semiclassical entropy induced by the quantum hair  $N$  and the semiclassical results are reproduced even when  $\beta$  does not exactly obey the relation (119) and the chemical potential is not exactly zero. Moreover, investigating more in detail the constraint (118) from this point of view could provide useful insights to derive (119) from physical considerations.

To conclude this discussion on the role of the Barbero-Immirzi parameter, we have seen that, since the introduction of the basic conceptual ideas of [14]-[16] and the seminal works of [18, 19], research on black hole entropy in LQG has been very active. We now have a fully  $SU(2)$  invariant description available of the model and we understand better the thermodynamical properties of the quantum IH system. This provided us with a set of different possible interpretations of the role played by  $\beta$  in the entropy calculation. Even if the original numerical constraint found in [19] is still a valid possibility, this is no longer the only one. The statement that the LQG calculation recovers the Bekenstein-Hawking area law *only* by fixing the Barbero-Immirzi parameter to a specific numerical value is therefore no longer true!

Surely, further study is necessary to understand better the semi-classical

limit of IH.

## 9 Derivation from Conformal Field Theories

In this section we are going to present a different point of view on the computation of entropy, paying special attention to the theoretical structure of the framework and the possible underlying symmetries that could take part in it, following the work in [34]. Motivated by the results of [11, 12], we want to search for the possible interplay between the theory describing black holes in loop quantum gravity and a possible underlying conformal symmetry.

By paying special attention to the fact that the horizon is described by a Chern-Simons theory, we are now going to make use of Witten's proposal about the connection between Chern-Simons theories and Wess-Zumino-Witten models. More precisely, in [65] Witten proposed the correspondence between the Hilbert space of generally covariant theories and the space of conformal blocks of a conformally invariant theory. This idea has been applied in [67] to the computation of the entropy for a horizon described by a  $SU(2)$  Chern-Simons theory, by putting its Hilbert space in correspondence with the space of conformal blocks of a  $SU(2)$ -Wess-Zumino-Witten (WZW) model. In this Section we are now going to make use of Witten's correspondence for the  $U(1)$  Chern-Simons theory describing the black hole horizon in LQG, according to the model presented in Section 5.1, and look for some hints on the role of CFT techniques in this framework.

Taking into account the fact that this  $U(1)$  group arises as the result of a geometric symmetry breaking from the  $SU(2)$  symmetry in the bulk, one can still make use of the well established correspondence between  $SU(2)$  Chern-Simons and Wess-Zumino-Witten theories. However, in this case it will be necessary to impose restrictions on the  $SU(2)$ -WZW model, as we will see, in order to implement the symmetry reduction. Through this procedure we expect to eventually reproduce the counting of the Hilbert space dimension of the  $U(1)$  Chern-Simons theory.

Let us begin by recalling the classical scenario and how the symmetry reduction takes place at this level. The geometry of the bulk is described by a  $SU(2)$  connection, whose restriction to the horizon  $H$  gives rise to a  $SU(2)$  connection over this surface. As a way of imposing the isolated horizon boundary conditions, this connection can be reduced to a  $U(1)$  connection. In [19] this reduction is carried out, at the classical level, just by fixing a unit vector  $\vec{r}$  at each point of the horizon. By defining a smooth function  $r : S \rightarrow \mathfrak{su}(2)$  a  $U(1)$  sub-bundle is picked out from the  $SU(2)$  bundle. This kind of reduction can be described in more general terms as follows (see, for instance, [80]). Let  $P(SU(2), S)$  be a  $SU(2)$  principal bundle over the horizon, and  $\omega$  the corresponding connection over it. A homomorphism  $\lambda$  between the

closed subgroup  $U(1) \subset SU(2)$  and  $SU(2)$  induces a bundle reduction from  $P(SU(2), S)$  to  $Q(U(1), S)$ ,  $Q$  being the resulting  $U(1)$  principal bundle with reduced  $U(1)$  connection  $\omega'$ . This  $\omega'$  is obtained, in this case, from the restriction of  $\omega$  to  $U(1)$ . All the conjugacy classes of homomorphisms  $\lambda : U(1) \rightarrow SU(2)$  are represented in the set  $Hom(U(1), T(SU(2)))$ , where  $T(SU(2)) = \{diag(z, z^{-1}) | z = e^{i\theta} \in U(1)\}$  is the maximal torus of  $SU(2)$ .

The homomorphisms in  $Hom(U(1), T(SU(2)))$  can be characterized by

$$\lambda_p : z \mapsto diag(z^p, z^{-p}) , \quad (120)$$

for any  $p \in \mathbb{Z}$ . However, the generator of the Weyl group of  $SU(2)$  acts on  $T(SU(2))$  by  $diag(z, z^{-1}) \mapsto diag(z^{-1}, z)$ . If we divide out by the action of the Weyl group we are just left with those maps  $\lambda_p$  with  $p$  a non-negative integer,  $p \in \mathbb{N}_0$ , as representatives of all conjugacy classes. These  $\lambda_p$  characterize then all the possible ways to carry out the symmetry breaking from the  $SU(2)$  to the  $U(1)$  connection that will be quantized later.

In alternative, one can follow another approach consisting of first quantizing the  $SU(2)$  connection on  $H$  and imposing the boundary conditions later on, at the quantum level. This would give rise to a  $SU(2)$  Chern-Simons theory on the horizon on which the boundary conditions now have to be imposed. The correspondence with conformal field theories can be used at this point to compute the dimension of the Hilbert space of the  $SU(2)$  Chern-Simons as the number of conformal blocks of the  $SU(2)$ -WZW model, as it was done in [67]. It is necessary to require, then, additional restrictions to the  $SU(2)$ -WZW model that account for the symmetry breaking, and consider only the degrees of freedom corresponding to a  $U(1)$  subgroup.

Let us briefly review the computation in the  $SU(2)$  case, to later introduce the symmetry reduction. The number  $N^{\mathcal{P}}$  of conformal blocks of the  $SU(2)$ -WZW model, given a set of representations  $\mathcal{P} = \{j_1, j_2, \dots, j_N\}$ , can be computed in terms of the so-called fusion numbers  $\mathcal{N}_{il}^r$  [81] as

$$N^{\mathcal{P}} = \sum_{r_i} \mathcal{N}_{j_1 j_2}^{r_1} \mathcal{N}_{r_1 j_3}^{r_2} \dots \mathcal{N}_{r_{N-2} j_{N-1}}^{j_N} . \quad (121)$$

These  $\mathcal{N}_{il}^r$  are the number of independent couplings between three primary fields, *i.e.* the multiplicity of the  $r$ -irreducible representation in the decomposition of the tensor product of the  $i$  and  $l$  representations  $[j_i] \otimes [j_l] = \bigoplus_r \mathcal{N}_{il}^r [j_r]$ . This expression is known as a fusion rule.  $N^{\mathcal{P}}$  is then the multiplicity of the  $SU(2)$  gauge invariant representation ( $j = 0$ ) in the direct sum decomposition of the tensor product  $\bigotimes_{i=1}^N [j_i]$  of the representations in  $\mathcal{P}$ . The usual way of computing  $N^{\mathcal{P}}$  is using the Verlinde formula [81] to obtain the fusion numbers. But alternatively one can make use of the fact that the characters of the  $SU(2)$  irreducible representations,  $\chi_i = \sin[(2j_i + 1)\theta] / \sin \theta$ , satisfy the fusion rule  $\chi_i \chi_j = \sum_r \mathcal{N}_{ij}^r \chi_r$ . Taking into account that the characters form an orthonormal set with respect to

the  $SU(2)$  scalar product,  $\langle \chi_i | \chi_j \rangle_{SU(2)} = \delta_{ij}$ , one can obtain the number of conformal blocks just by projecting the product of characters over the character  $\chi_0$  of the gauge invariant representation

$$N^{\mathcal{P}} = \langle \chi_{j_1} \dots \chi_{j_N} | \chi_0 \rangle_{SU(2)} = \int_0^{2\pi} \frac{d\theta}{\pi} \sin^2 \theta \prod_{i=1}^N \frac{\sin [(2j_i + 1)\theta]}{\sin \theta} . \quad (122)$$

This expression is equivalent to the one obtained in [67] using the Verlinde formula; it gives rise to the same result for every set of punctures  $\mathcal{P}$ .

To implement, now, the symmetry breaking we have to restrict the representations in  $\mathcal{P}$  to a set of  $U(1)$  representations. In the case of Chern-Simons theory, this corresponds to performing a symmetry reduction locally at each puncture. It is known that each  $SU(2)$  irreducible representation  $j$  contains the direct sum of  $2j + 1$   $U(1)$  representations  $e^{ij\theta} \oplus e^{i(j-1)\theta} \oplus \dots \oplus e^{-ij\theta}$ . One can make an explicit symmetry reduction by just choosing one of the possible restrictions of  $SU(2)$  to  $U(1)$  which, as we saw above, are given by the homomorphisms  $\lambda_p$ . This amounts here to pick out a  $U(1)$  representation of the form  $e^{ip\theta} \oplus e^{-ip\theta}$  with some  $p \leq j$ . The fact that we will be using these reducible representations, consisting of  $SU(2)$  elements as  $U(1)$  representatives, can be seen as a reminiscence from the fact that the  $U(1)$  freedom has its origin in the reduction from  $SU(2)$ .

Having implemented the symmetry reduction, let us compute the number of independent couplings in this  $U(1)$ -reduced case. Of course, we are considering now  $U(1)$  invariant couplings, so we have to compute the multiplicity of the  $m = 0$  irreducible  $U(1)$  representation in the direct sum decomposition of the tensor product of the representations involved. As in the previous case, this can be done by using the characters of the representations and the fusion rules they satisfy. These characters can be expressed as  $\tilde{\eta}_{p_i} = e^{ip_i\theta} + e^{-ip_i\theta} = 2 \cos p_i\theta$ . Again, we can make use of the fact that the characters  $\eta_i$  of the  $U(1)$  irreducible representations are orthonormal with respect to the standard scalar product in the circle. Then, the number we are looking for is given by

$$N_{U(1)}^{\mathcal{P}} = \langle \tilde{\eta}_{p_1} \dots \tilde{\eta}_{p_N} | \eta_H \rangle_{U(1)} = \frac{1}{2\pi} \int_0^{2\pi} d\theta \prod_i^N 2 \cos p_i\theta , \quad (123)$$

where  $\eta_H = 1$  is the character of the  $U(1)$  gauge invariant irreducible representation. We can see that this result is exactly the same as the one obtained for  $P(\{n_s\})$  in eq. (106), coming from the  $U(1)$  Chern-Simons theory, just by identifying the  $p_i$  with  $s_i$  labels.

From the physical point of view, the main change we are introducing, besides using the Chern-Simons/CFT analogy, is to impose the isolated horizon boundary conditions at the quantum level, instead of doing it prior to

the quantization process. This can be seen as a preliminary step in the direction of introducing a quantum definition of isolated horizons.

## 10 Observational Tests

We would like to end this review by commenting on some of the most recent results on black hole entropy.

We have seen that the combinatorial problem giving rise to the entropy counting is quite an elaborate one, and some somewhat technical steps are required to solve it. Furthermore, there is a nice interplay between the different particular structures involved at each step that gives rise to non-trivial structures on the degeneracy spectrum of black holes. In particular, the observed band structure for microscopic black holes is a very characteristic signature, and the precise features of the loop quantum gravity area spectrum play a major role in this result.

One can ask whether all this detailed structures could have an influence on some physical processes, like Hawking radiation, and whether they could give rise to observable effects. That possibility was already conjectured in [31], on the basis of a qualitative spectroscopical analysis. However, one could try to use some computational methods – in particular Monte Carlo simulations – to test if there is actually such an observational signal, and whether it would be possible to discriminate between loop quantum gravity and the standard semi-classical approach (or other quantum gravity theories) by observing microscopic black hole evaporation. This question was very recently addressed in [46]. In that work, a Monte Carlo simulation was performed, using precise data on the black hole degeneracy spectrum as an input. The transition probability between states was modulated by a factor proportional to the degeneracy of the final state. In particular, following [82], a factor of the form

$$P_{1 \rightarrow 2} = N e^{-\Delta S_{12}} \quad (124)$$

was introduced, where  $\Delta S_{12}$  is the difference in entropy between the initial and final states, and  $N$  is a gray-body factor, whose exact value was computed numerically. The radiation spectrum resulting from the Monte Carlo simulation is shown in Figure 6. Some characteristic lines superimposed to the quasi-continuous spectrum can be clearly appreciated. On the basis of this observation, a detailed statistical analysis was performed in order to determine whether that signal could be discriminated from the predictions of other black hole models. A Kolmogorov-Smirnov test – measuring the distance between the cumulative distribution functions of both distributions – was run to determine the ability to discriminate between loop quantum gravity and the semiclassical Hawking spectrum in an hypothetical observation. The results are shown in Figure 7, where the deviation between




Figure 6: Spectrum of emitted particles both for loop quantum gravity (up) and for the semiclassical Hawking case, as resulting from the Monte Carlo simulation.

models (for several confidence levels) is plotted as a function of the number of observed black holes and the relative error in the observation. It can be seen that, either a large enough number of observed black holes, or a small enough relative error, would allow to discriminate between both models. Additional tests comparing loop quantum gravity with other discrete models were also performed, showing an even better result in the discrimination. This shows that a (hypothetical) observation of microscopic black hole evaporation could be used for probing loop quantum gravity.




Figure 7: Number of observed evaporating black holes needed to discriminate loop quantum gravity and semiclassical Hawking models as a function of the relative error in the energy reconstruction of the emitted particles and the confidence level.

A remarkable fact about these considerations is that the specific signatures that are used to probe loop quantum gravity arise as a consequence, not so much of the particular model used for the black hole description, but of the structure of the area spectrum in the theory. This fact makes the resulting predictions much more robust, since it is reasonable to think that they are independent of the particular assumptions made for the current description of black holes in loop quantum gravity, and therefore they could be expected to remain valid – up to a certain extent – even after a full quantum description of a black hole is available in the theory.

Another possible observational test of the theory coming from the measurement of the Hawking radiation spectrum has been very recently conjectured in [61]. There, the authors propose a modification of the black hole radiation spectrum in relation to an additional term introduced in the first law and proportional to the variation of the number of punctures contributing to the macroscopic geometry of the horizon [45]. According to the usual matter coupling in LQG, one would expect the emission/absorption of fermions to induce a change in the number of punctures piercing the horizon and this process would, therefore, become observable if such a modification

of the first law affected the black hole radiation spectrum, as proposed by [61].

## 11 Conclusions

The quasi-local definition of black hole encoded in the notion of isolated horizon, for which the familiar laws continue to hold, provides a physically relevant and suitable framework to start a quantization program of the boundary degrees of freedom. By extracting an appropriate sector of the theory in which space-time geometries satisfy suitable conditions at an inner boundary representing the horizon – to ensure only that the intrinsic geometry of the horizon be time independent although the geometry outside may be dynamical and admit gravitational and other radiation –, one can construct the Hamiltonian framework and derive a conserved symplectic structure for the system. As shown in Section 3, when switching from the vector-like variables to the (Ashtekar-Barbero) connection variables in the bulk theory, in order to later allow the use of techniques developed for quantization, the symplectic form acquires a boundary contribution.

There is a certain freedom in the choice of boundary variables leading to different parametrizations of the boundary degrees of freedom. The most direct description would appear, at first sight, to be the one defined simply in terms of the triad field (pulled back on  $H$ ). Such a parametrization is however less preferable from the point of view of quantization, as one is confronted with the background independent quantization of form fields for which the usual techniques are not directly applicable; moreover, as discussed in Section 3.3, with this parametrization the entropy may be affected by the presence of degenerate geometry configurations left over after the imposition of the boundary constraint. In contrast, the parametrization of the boundary degrees of freedom in terms of connections directly leads to a description in terms of Chern-Simons theory which, being a well-studied topological field theory, drastically simplifies the problem of quantization. This allows us to obtain a remarkably simple formula for the horizon entropy: the number of states of the horizon is simply given in terms of the (well-known) dimension of the Hilbert spaces of Chern-Simons theory with punctures labeled by spins.

Performing a  $U(1)$  gauge fixing provides a classically equivalent description of the boundary degrees of freedom, but has some important implications in the quantum theory. One of these is the different numerical factor in front of the logarithmic corrections. Avoiding the gauge reduction preserves the full  $SU(2)$  nature of the IH quantum constraints, allowing us to impose them strongly in the Dirac sense. This leads to sub-leading corrections of the form  $\Delta S = -\frac{3}{2} \log a_H$ , matching other approaches [10, 12]. This way, the relevance of the original intuition of [67], which yielded a universal form of

logarithmic corrections in agreement with those found in different quantum gravity formulations, is clarified and put on solid ground.

However, the  $SU(2)$  invariant description comes with the freedom of the introduction of an extra dimensionless parameter. Such an appearance of extra parameters is intimately related to what happens in the general context of the canonical formulation of gravity in terms of connections. Therefore, this observation is by no means a new feature characteristic of IH. The existence of this extra parameter has a direct influence on the value of the Chern-Simons level. As shown in Section 6.2, this can be used to define an effective theory in which the entropy of the horizon grows as  $a_H/(4\ell_p^2)$  without the need of fixing the Barbero-Immirzi parameter to any specific value, but simply imposing a given relation between  $\beta$  and  $k$ . Gaining physical insight on this relation between (a parameter proper of) the bulk and the boundary theory may have potentially deep implications. In this direction, the new ‘relocation’ of the Barbero-Immirzi parameter in the entropy formula provided by the recent analysis of [45] could turn out to be very useful.

A remarkable fact is that, despite the various improvements and constant evolution of the framework, most of the powerful techniques developed in [35, 36, 37] for the entropy computation are still useful –with slight modifications– to solve the counting problem within this recently developed approach [43]. Furthermore, the effective quantization of entropy, resulting from the discrete nature of the problem, has proved to be a robust feature, appearing repeatedly regardless of the approach followed. With the recent analysis [38] showing the disappearance of this effect for large horizon areas, the entropy discretization remains as a robust prediction of this framework for black holes in the Planck regime. This consistency is particularly important to support the study of possible observational signatures arising as a consequence of the discretization effect.

The inclusion of distortion has been recently implemented, both in the  $U(1)$  and in the  $SU(2)$  formulation. In the former case, this has been performed by ‘reinterpreting’ the original Hilbert space of [19], through the mapping to a fiducial Type I structure, as the quantum counterpart of the full phase-space of all distorted IH of a given area [76]. In the latter case, horizon distortion can be taken into account by introducing two  $SU(2)$  Chern-Simons theories on the boundary [42]. This allows to define a quantum operator encoding the distortion degrees of freedom, whose eigenvalues are expressed in terms of the spins associated to the bulk and the boundary punctures.

A better understanding of the relation between these two pictures could be provided by a characterization of the horizon theory from the full theory. The first steps in this direction have already been moved in [63, 64]. Here the authors start from the flux-holonomy algebra of LQG which represents a quantization of the kinematical degrees of freedom of GR in the connec-

tion formulation. Studying a modification of the Ashtekar-Lewandowski measure on the space of generalized connections, one can look for a representation of this algebra containing states that solve the quantum analog the boundary conditions and thus provide a quantum mechanics description of black holes. Therefore, the approach taken in [63, 64] differs from the one adopted in [19, 41], since there the boundary and bulk degrees of freedom are no longer treated separately at the quantum level. On the contrary, the horizon degrees of freedom are now represented simply by elements of the flux-holonomy algebra of LQG, without any reference to the horizon. Providing a characterization of the operators entering the IH boundary conditions from the full quantum theory definitely represents a very important step and it might provide a deeper understanding of the horizon quantum geometry degrees of freedom, with the possibility to give new insights on the relation between the models defined in [76] and [42].

Finally, the recent study showing that the particular features of loop quantum gravity produce observational signatures that are relevant enough to allow a clear discrimination between this theory and other possible quantum black hole models on a simulated experiment [46], is a strong encouragement to keep extending and improving the understanding of quantum black holes, and to tackle with interest the remaining open issues. After all, who knows if they could be the key to the first observational test of quantum gravity?

## Acknowledgements

We would like to specially thank Abhay Ashtekar and Alejandro Perez. This work was partially supported by NSF grants PHY-0854743 and PHY-0968871, the Spanish MICINN grant ESP2007-66542-C04-01, and the Eberly research funds of Penn State.

## References

- [1] M. J. Reid, “Is there a Supermassive Black Hole at the Center of the Milky Way?,” *Int. J. Mod. Phys. D* **18** (2009) 889 [arXiv:0808.2624 [astro-ph]].  
A. Mueller, “Experimental evidence of black holes,” *PoS P2GC* (2006) 017 [arXiv:astro-ph/0701228].  
A. E. Broderick, A. Loeb and R. Narayan, “The Event Horizon of Sagittarius A\*,” *Astrophys. J.* **701** (2009) 1357 [arXiv:0903.1105 [astro-ph.HE]].
- [2] R. M. Wald, “General Relativity,” *Chicago, Usa: Univ. Pr. (1984) 491p.*

- [3] R. P. Kerr, “Gravitational field of a spinning mass as an example of algebraically special metrics” *Phys. Rev. Lett.* **11** (1963) 237-238.  
E. T. Newman, R. Couch, K. Chinnapared, A. Exton, A. Prakash and R. Torrence, “Metric of a Rotating, Charged Mass,” *J. Math. Phys.* **6** (1965) 918.
- [4] S. W. Hawking and G. F. R. Ellis, “The Large scale structure of space-time,” *Cambridge University Press, Cambridge, 1973*.
- [5] M. Bojowald, “Nonsingular black holes and degrees of freedom in quantum gravity,” *Phys. Rev. Lett.* **95**, 061301 (2005) [arXiv:gr-qc/0506128].
- [6] A. Ashtekar and M. Bojowald, “Black hole evaporation: A paradigm,” *Class. Quant. Grav.* **22**, 3349 (2005) [arXiv:gr-qc/0504029].
- [7] A. Ashtekar and M. Bojowald, “Quantum geometry and the Schwarzschild singularity,” *Class. Quant. Grav.* **23**, 391 (2006) [arXiv:gr-qc/0509075].
- [8] J. D. Bekenstein, “Black holes and entropy,” *Phys. Rev. D* **7** (1973) 2333.
- [9] S. W. Hawking, “Particle Creation By Black Holes,” *Commun. Math. Phys.* **43** (1975) 199 [Erratum-ibid. **46** (1976) 206].
- [10] A. Strominger and C. Vafa, “Microscopic Origin of the Bekenstein-Hawking Entropy,” *Phys. Lett. B* **379** (1996) 99 [arXiv:hep-th/9601029].
- [11] A. Strominger, “Black hole entropy from near horizon microstates,” *JHEP* **9802**, 009 (1998). [hep-th/9712251].
- [12] S. Carlip, “Entropy from conformal field theory at Killing horizons,” *Class. Quant. Grav.* **16** (1999) 3327 [arXiv:gr-qc/9906126].  
S. Carlip, “Black hole entropy from conformal field theory in any dimension,” *Phys. Rev. Lett.* **82** (1999) 2828 [arXiv:hep-th/9812013].  
S. Carlip, “Black Hole Thermodynamics and Statistical Mechanics,” *Lect. Notes Phys.* **769** (2009) 89 [arXiv:gr-qc/0807.4520].
- [13] A. Ashtekar and J. Lewandowski, “Background independent quantum gravity: A status report,” *Class. Quant. Grav.* **21**, R53 (2004) [arXiv:gr-qc/0404018].  
C. Rovelli, “Quantum Gravity,” *Cambridge, UK: Univ. Pr. (2004) 455 p.*  
T. Thiemann, “Modern canonical quantum general relativity,” *Cambridge, UK: Univ. Pr. (2007) 819 p.*

- A. Perez, "Introduction to loop quantum gravity and spin foams," [arXiv:gr-qc/0409061].
- [14] L. Smolin, "Linking topological quantum field theory and nonperturbative quantum gravity," *J. Math. Phys.* **36** (1995) 6417 [arXiv:gr-qc/9505028].
- [15] C. Rovelli, "Black hole entropy from loop quantum gravity," *Phys. Rev. Lett.* **77** (1996) 3288 [arXiv:gr-qc/9603063].
- [16] K. V. Krasnov, "On statistical mechanics of gravitational systems," *Gen. Rel. Grav.* **30** (1998) 53 [arXiv:gr-qc/9605047].
- [17] A. Ashtekar, J. Baez, A. Corichi, K. Krasnov, "Quantum geometry and black hole entropy," *Phys. Rev. Lett.* **80**, 904-907 (1998). [gr-qc/9710007].
- [18] A. Ashtekar, A. Corichi and K. Krasnov, "Isolated horizons: The classical phase-space," *Adv. Theor. Math. Phys.* **3** (2000) 419 [arXiv:gr-qc/9905089].
- [19] A. Ashtekar, J. C. Baez, K. Krasnov, "Quantum geometry of isolated horizons and black hole entropy," *Adv. Theor. Math. Phys.* **4**, 1-94 (2000). [gr-qc/0005126].
- [20] A. Ashtekar, V. Taveras, M. Varadarajan, "Information is Not Lost in the Evaporation of 2-dimensional Black Holes," *Phys. Rev. Lett.* **100** (2008) 211302. [arXiv:0801.1811 [Unknown]].
- [21] A. Ashtekar, C. Beetle, S. Fairhurst, "Isolated horizons: A Generalization of black hole mechanics," *Class. Quant. Grav.* **16**, L1-L7 (1999). [gr-qc/9812065].
- [22] S. A. Hayward, "General laws of black hole dynamics," *Phys. Rev.* **D49**, 6467-6474 (1994).  
S. A. Hayward, "Spin coefficient form of the new laws of black hole dynamics," *Class. Quant. Grav.* **11**, 3025-3036 (1994). [gr-qc/9406033].
- [23] A. Ashtekar, B. Krishnan, "Isolated and dynamical horizons and their applications," *Living Rev. Rel.* **7**, 10 (2004). [gr-qc/0407042].
- [24] I. Booth, "Black hole boundaries," *Can. J. Phys.* **83**, 1073-1099 (2005). [gr-qc/0508107].
- [25] M. Domagala and J. Lewandowski, "Black hole entropy from quantum geometry," *Class. Quant. Grav.* **21** (2004) 5233 [arXiv:gr-qc/0407051].
- [26] K. A. Meissner, "Black hole entropy in loop quantum gravity," *Class. Quant. Grav.* **21** (2004) 5245 [arXiv:gr-qc/0407052].

- [27] G. Gour, “Algebraic approach to quantum black holes: Logarithmic corrections to black hole entropy,” *Phys. Rev.* **D66**, 104022 (2002). [gr-qc/0210024].
- A. Ghosh and P. Mitra, “A bound on the log correction to the black hole area law,” *Phys. Rev. D* **71** (2005) 027502 [arXiv:gr-qc/0401070].
- [28] A. Ghosh and P. Mitra, “An improved lower bound on black hole entropy in the quantum geometry approach,” *Phys. Lett. B* **616** (2005) 114 [arXiv:gr-qc/0411035].
- A. Ghosh and P. Mitra, “Counting black hole microscopic states in loop quantum gravity,” *Phys. Rev. D* **74** (2006) 064026 [arXiv:hep-th/0605125].
- A. Ghosh and P. Mitra, “Fine-grained state counting for black holes in loop quantum gravity,” *Phys. Rev. Lett.* **102** (2009) 141302 [arXiv:0809.4170 [gr-qc]].
- [29] A. Corichi, J. Diaz-Polo, E. Fernandez-Borja, “Quantum geometry and microscopic black hole entropy,” *Class. Quant. Grav.* **24**, 243-251 (2007). [gr-qc/0605014].
- [30] A. Corichi, J. Diaz-Polo, E. Fernandez-Borja, “Black hole entropy quantization,” *Phys. Rev. Lett.* **98**, 181301 (2007). [gr-qc/0609122].
- [31] J. Diaz-Polo and E. Fernandez-Borja, “Black hole radiation spectrum in loop quantum gravity: isolated horizon framework,” *Class. Quantum Grav.* **25**, 105007 (2008). [arXiv:0706.1979[gr-qc]].
- [32] H. Sahlmann, “Toward explaining black hole entropy quantization in loop quantum gravity,” *Phys. Rev. D* **76** (2007) 104050 [arXiv:0709.2433 [gr-qc]].
- H. Sahlmann, “Entropy calculation for a toy black hole,” *Class. Quant. Grav.* **25** (2008) 055004 [arXiv:0709.0076 [gr-qc]].
- [33] A. Agullo, J. Diaz-Polo and E. Fernandez-Borja “Black hole state degeneracy in Loop Quantum Gravity,” *Phys. Rev.* **D77** (2008) 104024. [arXiv:0802.3188[gr-qc]].
- [34] I. Agullo, J. Diaz-Polo and E. Fernandez-Borja, “Computing black hole entropy in loop quantum gravity from a conformal field theory perspective”, *JCAP* **0907**, 016 (2009), [arXiv:0903.1667 [hep-th]].
- [35] I. Agullo, J. F. Barbero G., J. Diaz-Polo, E. Fernandez-Borja and E. J. S. Villasenor, “Black hole state counting in LQG: A number theoretical approach”, *Phys. Rev. Lett.* **100** (2008) 211301; [arXiv:gr-qc/0802.4077].

- [36] J. F. Barbero G. and E. J. S. Villasenor, “Generating functions for black hole entropy in Loop Quantum Gravity”, *Phys. Rev. D* **77** (2008) 121502; [arXiv:gr-qc/0804.4784].  
 J. F. Barbero G. and E. J. S. Villasenor, “On the computation of black hole entropy in loop quantum gravity”, *Class. Quant. Grav.* **26** (2009) 035017; [arXiv:gr-qc/0810.1599].
- [37] I. Agullo, J. Fernando Barbero, E. F. Borja, J. Diaz-Polo and E. J. S. Villasenor, “Detailed black hole state counting in loop quantum gravity,” *Phys. Rev. D* **82** (2010) 084029.
- [38] G. J. Fernando Barbero, E. J. S. Villasenor, “Statistical description of the black hole degeneracy spectrum,” *Phys. Rev.* **D83**, 104013 (2011). [arXiv:1101.3662 [gr-qc]].
- [39] K. Krasnov, C. Rovelli, “Black holes in full quantum gravity,” *Class. Quant. Grav.* **26**, 245009 (2009). [arXiv:0905.4916 [gr-qc]].
- [40] J. Engle, A. Perez, K. Noui, “Black hole entropy and SU(2) Chern-Simons theory,” *Phys. Rev. Lett.* **105**, 031302 (2010). [arXiv:0905.3168 [gr-qc]].
- [41] J. Engle, K. Noui, A. Perez, D. Pranzetti, “Black hole entropy from an SU(2)-invariant formulation of Type I isolated horizons,” *Phys. Rev.* **D82**, 044050 (2010). [arXiv:1006.0634 [gr-qc]].
- [42] A. Perez, D. Pranzetti, “Static isolated horizons: SU(2) invariant phase-space, quantization, and black hole entropy,” *Entropy* 2011, 13(4), 744-777. [arXiv:1011.2961 [gr-qc]].
- [43] I. Agullo, J. F. Barbero G., E. F. Borja, J. Diaz-Polo and E. J. S. Villasenor, “The combinatorics of the SU(2) black hole entropy in loop quantum gravity”, *Phys. Rev. D* **80** (2009) 084006; [arXiv:gr-qc/0906.4529].
- [44] J. Engle, K. Noui, A. Perez, D. Pranzetti, “The SU(2) Black Hole entropy revisited,” *JHEP* **1105**, 016 (2011). [arXiv:1103.2723 [gr-qc]].
- [45] A. Ghosh, A. Perez, “Black hole entropy and isolated horizons thermodynamics,” [arXiv:1107.1320 [gr-qc]].
- [46] A. Barrau, T. Cailleteau, X. Cao, J. Diaz-Polo and J. Grain, “Probing Loop Quantum Gravity with Evaporating Black Holes,” [arXiv:1109.4239[gr-qc]].
- [47] A. Ashtekar, C. Beetle, S. Fairhurst, “Mechanics of isolated horizons,” *Class. Quant. Grav.* **17**, 253-298 (2000). [gr-qc/9907068].

- [48] A. Ashtekar, S. Fairhurst, B. Krishnan, “Isolated horizons: Hamiltonian evolution and the first law,” *Phys. Rev.* **D62**, 104025 (2000). [gr-qc/0005083].
- [49] A. Ashtekar, C. Beetle, O. Dreyer, S. Fairhurst, B. Krishnan, J. Lewandowski, J. Wisniewski, “Isolated horizons and their applications,” *Phys. Rev. Lett.* **85**, 3564-3567 (2000). [gr-qc/0006006].
- [50] A. Ashtekar, C. Beetle, J. Lewandowski, “Mechanics of rotating isolated horizons,” *Phys. Rev.* **D64**, 044016 (2001). [gr-qc/0103026].
- [51] A. Ashtekar, C. Beetle, J. Lewandowski, “Geometry of generic isolated horizons,” *Class. Quant. Grav.* **19**, 1195-1225 (2002). [gr-qc/0111067].
- [52] J. Lewandowski, “Space-times admitting isolated horizons,” *Class. Quant. Grav.* **17**, L53-L59 (2000). [gr-qc/9907058].
- [53] A. Rendall, “Reduction of the characteristic initial value problem to the Cauchy problem and its applications to Einstein’s equations,” *Proc. R. Soc.* **427** 221-239 (1990).
- [54] A. Ashtekar, J. Engle, T. Pawlowski, C. Van Den Broeck, “Multipole moments of isolated horizons,” *Class. Quant. Grav.* **21**, 2549-2570 (2004). [gr-qc/0401114].  
A. Ashtekar, J. Engle and C. Van Den Broeck, “Quantum horizons and black hole entropy: Inclusion of distortion and rotation,” *Class. Quant. Grav.* **22** (2005) L27 [arXiv:gr-qc/0412003].
- [55] S. Chandrasekhar, “The mathematical theory of black holes,” *Oxford, UK: Clarendon (1992) 646 p.*
- [56] R.P. Geroch, A. Held and R. Penrose, “A space-time calculus based on pairs of null directions,” *J. Math. Phys.* **1973**, 14, 874.
- [57] C. Crnkovic and E. Witten, “Covariant description of canonical formalism in geometrical theories,” in *Three Hundred Years of Gravitation*, Print-86-1309 (PRINCETON); Hawking, S.W., Israel, W., Eds.; Cambridge University Press: Cambridge, UK, 1989.
- [58] J. Lee and R.M. Wald, “Local symmetries and constraint,” *J. Math. Phys.* **1990**, 31, 725.
- [59] A. Corichi, E. Wilson-Ewing, “Surface terms, Asymptotics and Thermodynamics of the Holst Action,” *Class. Quant. Grav.* **27**, 205015 (2010). [arXiv:1005.3298 [gr-qc]].

- [60] Lewandowski, J.; Okolow, A.; Sahlmann, H.; Thiemann, T. “Uniqueness of the diffeomorphism invariant state on the quantum holonomy-flux algebra,” *Commun. Math. Phys.* **2006**, 267, 703–733. [arXiv:gr-qc/0504147].
- C. Fleischhack, “Representations of the Weyl Algebra in Quantum Geometry,” *Commun. Math. Phys.* **285** (2009) 67. [arXiv:math-ph/0407006].
- [61] E. Frodden, A. Ghosh, A. Perez, “A local first law for isolated horizons,” [arXiv:1110.4055 [gr-qc]].
- [62] K. V. Krasnov, “Quantum geometry and thermal radiation from black holes,” *Class. Quant. Grav.* **16**, 563-578 (1999). [gr-qc/9710006].
- G. J. F. Barbero, E. J. S. Villasenor, “The thermodynamic limit and black hole entropy in the area ensemble,” *Class. Quant. Grav.* **28**, 215014 (2011). [arXiv:1106.3179 [gr-qc]].
- [63] H. Sahlmann, “Black hole horizons from within loop quantum gravity,” *Phys. Rev.* **D84**, 044049 (2011). [arXiv:1104.4691 [gr-qc]].
- [64] H. Sahlmann and T. Thiemann, “Chern-Simons expectation values and quantum horizons from LQG and the Duffo map” [arXiv:1109.5793[gr-qc]].
- [65] E. Witten, “Quantum field theory and the Jones polynomial,” *Commun. Math. Phys.* **121** (1989) 351.
- [66] T. Thiemann, “Quantum spin dynamics. VIII. The Master constraint,” *Class. Quant. Grav.* **23**, 2249-2266 (2006). [gr-qc/0510011].
- T. Thiemann, “The Phoenix project: Master constraint program for loop quantum gravity,” *Class. Quant. Grav.* **23**, 2211-2248 (2006). [arXiv:gr-qc/0305080 [gr-qc]].
- [67] R. K. Kaul, P. Majumdar, “Quantum black hole entropy,” *Phys. Lett.* **B439**, 267-270 (1998). [gr-qc/9801080].
- R. K. Kaul, P. Majumdar, “Logarithmic correction to the Bekenstein-Hawking entropy,” *Phys. Rev. Lett.* **84**, 5255-5257 (2000). [gr-qc/0002040].
- S. Das, R. K. Kaul, P. Majumdar, “A New holographic entropy bound from quantum geometry,” *Phys. Rev.* **D63**, 044019 (2001). [hep-th/0006211].
- [68] S. Carlip, “Logarithmic corrections to black hole entropy from the Cardy formula,” *Class. Quant. Grav.* **17** (2000) 4175 [arXiv:gr-qc/0005017].

- [69] E. Bianchi, “Black Hole Entropy, Loop Gravity, and Polymer Physics,” *Class. Quant. Grav.* **28**, 114006 (2011). [arXiv:1011.5628 [gr-qc]].
- [70] V. Chary, A. Pressley, “A guide to quantum groups,” *Cambridge University Press* (1994).
- [71] E. R. Livine, D. R. Terno, “Quantum black holes: Entropy and entanglement on the horizon,” *Nucl. Phys.* **B741**, 131-161 (2006). [gr-qc/0508085].  
 L. Freidel and E. R. Livine, “The Fine Structure of SU(2) Intertwiners from U(N) Representations,” *J. Math. Phys.* **51** (2010) 082502 [arXiv:0911.3553 [gr-qc]].  
 E. R. Livine and D. R. Terno, “The entropic boundary law in BF theory,” *Nucl. Phys. B* **806** (2009) 715 [arXiv:0805.2536 [gr-qc]].  
 E. R. Livine and D. R. Terno, “Bulk Entropy in Loop Quantum Gravity,” *Nucl. Phys. B* **794** (2008) 138 [arXiv:0706.0985 [gr-qc]].
- [72] D. J. Newman, W. F. Furr and L. K. Williams, *The American Mathematical Monthly* **66**, 321 (1959).
- [73] D. M. Burton, *Elementary Number Theory*, McGraw-Hill, New York (2002).
- [74] H. De Raedt, K. Michielsen, K. De Raedt and S. Miyashita, “Number partitioning on a quantum computer,” *Phys. Lett.* **A290**, 227-233 (2001). [arXiv:quant-ph/0010018]
- [75] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, Cambridge (2009).
- [76] C. Beetle, J. Engle, “Generic isolated horizons in loop quantum gravity,” *Class. Quant. Grav.* **27**, 235024 (2010). [arXiv:1007.2768 [gr-qc]].
- [77] C. Rovelli, T. Thiemann, “The Immirzi parameter in quantum general relativity,” *Phys. Rev.* **D57**, 1009-1014 (1998). [gr-qc/9705059].
- [78] R. Durka, J. Kowalski-Glikman, “Gravity as a constrained BF theory: Noether charges and Immirzi parameter,” *Phys. Rev.* **D83**, 124011 (2011). [arXiv:1103.2971 [gr-qc]].
- [79] R. M. Wald, “Black hole entropy is the Noether charge,” *Phys. Rev.* **D48**, 3427 (1993). [arXiv:gr-qc/9307038].
- [80] M. Bojowald, H. A. Kastrup, “Quantum symmetry reduction for diffeomorphism invariant theories of connections,” *Class. Quant. Grav.* **17**, 3009-3043 (2000). [hep-th/9907042].

- [81] P. Di Francesco, P. Mathieu and D. Senechal, *Conformal Field Theory*, Springer, New York(1997).
- [82] S. Massar and R. Parentani, “How the change in horizon area drives black hole evaporation”, Nucl. Phys. B, **575**, 333 (2000) [arXiv:gr-qc/9903027].